

Review

A review of the control of reproduction and hormonal manipulations in finfish species

Yousefian Mehdi* and Mousavi, Seyed Ehsan

Islamic Azad University, Qaemshahr branch, Iran.

Accepted 15 May, 2019

Control of reproductive is essential for the sustainability of commercial aquaculture production. The reason for breakage of ovulation and spawning is failure of the pituitary to release gonadotropin hormones. Gonadotropin hormones secretion is under a dual control, by gonadotropin releasing hormone (GnRH), and inhibition by dopamine on the actions of GnRH. The inhibitory actions of dopamine on GTH secretion can vary in potency between species. In some fish it is necessary to use dopamine blocker, such as domperidone, results in potentiation of the actions of GnRH analoge, leading to a large release of GTH-II and ovulation. In some other injection of a high dose of GnRH analoge alone is effective. In several marine species GTH-II secretion is not under dopaminergic inhibitory control but sensitive to exogenous analoge hormones (GnRH_a). In many fishes, synthetic analoges of lutenizing hormone-releasing hormone (LHRH_a) are widely used for inducing spawning. In several tropical freshwater and marine finfishes the human chorionic gonadotropin (HCG), pituitary homogenate and semi-purified fish gonadotropin are used. Rrecently the replacement of natural GnRH are used and had resulted in a centuple increase in the effectiveness of luteinizing hormones secretion induction.

Key words: Reproduction induced spawning, dopamine antagonists, GTH-II- HCG.

INTRODUCTION

The further development of aquaculture is undoubtedly based on the management of fish species. Aquaculture has been practiced for many centuries, but transfer from low and temporary production to intensive and regular approach is to control reproductive processes of fish in captivity, and to acquire high quality fingerlings for grow-out of the marketable product. Most of the fishes when reared in captivity, do not exhibit normal reproduction. Actually they reach reproduction maturity and gonadal growth is normal but the females fail final oocyte maturation (FOM) and do not spawn and male exhibit few or even lack of milt. The example for this subject is female common carp that in their habitat they will reproduce just by change in environmental condition and the presence of male. Therefore, breeding of captive fish may be approached managing the proper environmental condition suitable for the species. Artificial environments

lack natural spawning stimuli (spawning substrate, stream hydraulic, nutrition, water quality, depth etc.) are not able to induce appropriate endogenous response from the fish; the final result is reproductive dysfunction of FOM (Abraham, 1988; Podhorec and kouril, 2009).

Traditional methods of induced spawning for cultured fish are based on the injection of GTH-II from different sources, including crude extract of carp pituitary gland (CPE), partially purified fish GTH-II and mammalian GTH, especially human chorionic hormone (HCG) (Lam, 1982; Donaldson and Hunter, 1983; Peter et al., 1988a). Among these hormones, Fish GTH-II is generally of high species specificity, thus, for example, carp or salmon GTH-II are ineffective in gilthead seabream. HCG is routinely used to ineffective in others, such as grass carp and black carp (Lin and Peter, 1996). In using hormones for artificial reproduction it should be considered that several items such as the side effects of mixture of hormones, the cost of some hormones, also at the time of treatment, the fish should be at an advanced stage of gonadal development otherwise hormone injection at improper of time or

*Corresponding author. E-mail: yousefianeco@yahoo.com.

insufficient of the dosage may damage or even will kill the fish. Almost in all cultured fishes, hormonal manipulations may be used as management tools to enhance the efficiency of egg production, increase spermiation and facilitate hatchery operations. Also, hormonal therapies may be employed in artificial collection in order to implement interspecific hybridization, chromosome set manipulation or artificial fertilization for genetic selection programmes (Mylonas et al., 2010).

To have a review of the reproduction control and hormonal manipulations in finfish species, we may first have an investigation of the general hormonal function in reproduction and then the use of different hormone in different fish species. The objective of the present study was a brief study of different hormones used in fish reproduction and listing that are used in different fish species.

THE REPRODUCTIVE CYCLES IN FISH

The endocrine control system of the reproduction in finfish is based on the: hypothalamus-hypophysis-gonads axis. Therefore there is a permanent communication between the brain/pituitary complex and the periphery of the organ (gonads). This communication allows the activity of the different components of the brain-pituitary-gonad axis to be synchronized at all steps of the life cycle, which is crucial for coordinated responses. The rule of GnRH, secretion of dopamine, steroid feed back, LH secretion all has their importance alone or in combination with the others in reproduction. Two types of GtHs, different for structure and chemical role, were identified in the teleosts:

GtH-I (or FSH), involved in the initial stages of gametogenesis (vitellogenesis and spermatogenesis), and GtH-II (or LH), which sets FOM, spermiogenesis and spermiation (Mylonas et al., 1996).

GtH-II increase in plasma just before spermiation and FOM, determining the switch from the steroidogenic and androgenic production to the progestinic production respectively in the interstitial testicular cells and the ovarian theca cells (Rainis and Ballestrazzi, 2005). Studies comparing the relative LH releasing effects of the native GnRH isoforms in teleost species, including goldfish, catfish and seabream demonstrated that the most potent GnRH form in terms of bioactivity is GnRH-2, while the species-specific type 1 isoform is generally the least potent (Johnson et al., 1999; Rebers et al., 2000; Zohar et al., 1995b). In many teleosts, but not all, dopamine was shown to strongly inhibit gonadotropin release through a mechanism that probably has different adaptative significance depending on the species.

Dopamine is a small neurotransmitter that is synthesized from tyrosine through a two step reaction involving the step limiting enzyme tyrosine hydroxylase and DOPA-decarboxylase.

The distribution of dopamine in the brain of fish has been extensively studied using different techniques showing the existence of a well developed dopaminergic system (Nieuwenhuys et al., 1998). Dopamine receptors belong to the G-protein coupled receptor (GPCR) family. There are two main classes of DA receptors that differ in their ability to activate (D1) or to inhibit (D2) the enzyme adenylyl cyclase, with each class containing various subtypes (Cardinaud et al., 1998; Kebabian and Calne, 1979; Zohar et al., 2010). It was shown that dopamine acts directly onto the gonadotrophs through D2 receptors. This was shown first in goldfish and then confirmed in other species such as carp, African catfish, trout, tilapia, eel and gray mullet. In contrast, dopamine inhibition of gonadotropin release was not observed in many other teleost fish, particularly in marine species (see for more detail (Zohar et al., 2010). In fish species as mentioned before the rule of LH is very important in final maturation and ovulation of the egg. The neurodecapeptide GnRH is the central regulator of the reproductive hormonal cascade regulating the synthesis and release of LH secretion from the pituitary gland (Kah et al., 2007). The hypophysiotropic GnRH is processed in the hypothalamic neurons by enzymatic cleavage of a precursor polypeptide and packaged in storage granules (Yaron and Sivan, 2006). Another important factor controlling the reproduction in fish is the secretion and the rule of catecholamine metabolism. The teleost species possess two distinct gonadotropins, GtH-I and GtH-II. This is the case of seabream, striped bass, salmon and trout, whereas only one gonadotropin could be found in catfish (Saligaut et al., 1999).

GtH-I and GtH-II have distinct temporal expression and release profile in teleosts. GtH-I is released over the entire vitellogenesis, whereas GtH-II remains low during vitellogenesis and exhibit a sharp peak before ovulation (Davies et al., 1995; Prat et al., 1996). These different patterns of expression and release demonstrate that there is obviously a differential control of GtH-I and GtH-II (Saligaut et al., 1999). Studying more detail of the effect of GTH showed that GTH-I and GTH-II are equipotent in stimulating 11-ketotestosterone (11KT) and $17\alpha, 20\beta$ P production (Planas et al., 1991). By further study it was shown that later, however, GTH-II was more potent than GTH-I in stimulating $17\alpha, 20\beta$ P synthesis. Thus, early GnRH treatment may have had a relatively greater impact on GTH-I production while the effectiveness of GnRH_a injection at the height of the spawning season is consistent with increase impact on GTH-II production and $17\alpha, 20\beta$ P synthesis at that stage of the reproduction cycle. One of the primary factors limiting the early synthesis of $17\alpha, 20\beta$ P may also be the lack in availability of its precursor 17α -hydroxyprogesterone 17P

(Sakai et al., 1989; Schulz et al., 1991, 1992) due to the dominance of the androgen biosynthetic pathway. In salmonids, the role of catecholamines remains controversial. In salmon, DA modulated GtH release only after administration of a LHRH analogue (Van der Kraak et al., 1986). However, Billard et al. (1983) demonstrated in trout that pimozone (a DA antagonist) alone could increase plasma GtH.

The efficacy of pimozone and of MPT depends on E2 levels. In immature fish, the increase of GtH-II release is only observed in E2 implanted fish and occurs simultaneously with an increase of the activity of pituitary DA neurones (Linard et al., 1995). In both goldfish and trout, some dopaminergic neurones of the anterior ventral preoptic region project to proximal pars distalis of pituitary (Peter and Paulencu, 1980; Kah et al., 1987; Linard et al., 1996). An activation of a DA preoptico-hypophysial pathway could be involved during vitellogenesis in the rainbow trout. Some interesting results concern the catfish (*Heteropneustes fossilis*). It seems that in some air breathing species which coexist in sub-tropical waters, there is seasonality in the dominance of the CA during the reproductive cycle:

DA levels and turnover are high during the resting phase and decrease during the progress of breeding, with a concomitant increase in NE turnover (Senthikumar and Joy, 1995; Saligaut et al., 1999).

Oocyte maturation and spermiation

Blood levels of GtH-I in immature and vitellogenic trout are much higher than those of GtH-II and can be compared with those found in literature in salmon (Dickey and Swanson, 1995). In contrast GtH-II is high at final stage of gonad development. During vitellogenesis the oocyte development is continued. In investigation of "stock identification" and "genetic variation" at Microsatellite Loci of Caspian Sea Salmon (*Salmo trutta caspius*), genetic differentiation between the male and female, showed to be significant as indicated by Fst analysis, while the pair wise analysis of allele frequencies indicate differentiated at several loci, however we did not find the differences in sex is related to sex genes. (Yousefian, 2010b), at end of vitellogenesis oocyte maturation will be complete but for ovulation a number of cytological and nuclear changes occur. During FOM, the nucleus or germinal vesicle (GV) migrates to the periphery of the oocyte just under the micropyle. During or soon after GV migration, coalescence of the lipid droplets and yolk globules occurs, followed by the breakdown of the GV membrane (GVBD). This is the case that we usually use for testing the stage of artificially insemination of sturgeon fish in reproduction farms. In inducing artificial reproduction some important note should be considered.

The type of strategy employed for oogenesis by different fishes has important implications in their broodstock management in aquaculture. For example, for synchronous fishes that spawn once in their lifetime (for example, freshwater eels and Pacific salmon), sacrificing the fish and collecting their artificially matured gametes directly from the brood fish is a very effective and efficient method for producing fertilized eggs. Obviously, the same approach could not be used for fish that reach reproductive maturation at a late age and are expected to have a long reproductive life (for example, tunas, amberjacks or groupers). Similarly, stripping of eggs and artificial insemination may be used in annually spawning synchronous females to obtain the total amount of available eggs from each female, but the same may not be achieved with multiple spawning fishes that mature and ovulate only a small fraction of their total annual fecundity at every spawning event. During OM, some morphological changes occur in the oocyte together with progression of meiosis. In some such as mullet lipid droplet coalescence and yolk globule coalescence which result in the clarification of the oocyte's cytoplasm, migration of the nucleus (germinal vesicle, GV) to the periphery of the oocyte and dissolution of the nucleus membrane (GV breakdown, GVBD), and the volume of eggs increase due to water uptake. The GV is visible under the microscope after some chemical processing, and disappears when GVBD takes place (Yousefian et al., 2010a).

The problems in reproduction are not restricted to female but also low spermiation or low quality of sperm also may occur during spawning of fish. Lower plasma levels of LH during the spermiation period have been suggested as the cause of the reduced amount of milt produced by some fishes (Mananos et al., 2002; Mylonas and Zohar, 2001a). The amount of LH in the pituitary or the ability of the pituitary to synthesize LH in response to treatment with exogenous GnRHa is not affected in these fishes, suggesting that again the reproductive dysfunction in the males may be identified in the brain control of GtH synthesis and/or release (Mylonas et al., 2010). In case of female as we mentioned before the problem in reproduction is not due to vitellogenesis but the fish fail to undergo oocyte maturation. Steven and his colleagues in 2000, stated this problem that LH was synthesized and stored in the pituitary during vitellogenesis, since levels of LH and its mRNA in the pituitary did not differ between wild and captive females, demonstrating that the problem is one of lack of release and not synthesis in captivity. In addition, mRNA levels of the pituitary receptor for the GnRH most relevant to pituitary LH synthesis were similar between wild and captive females. This suggests that the disruption in LH release from the pituitaries of captive fish is not due to a dysfunction in pituitary responsiveness, but may be related to the control of pituitary function by reproductive brain. In fact, differences were observed between wild and captive

females undergoing OM, when comparing the pituitary content of the endogenous GnRHs.

The GnRH mRNA levels within the brain, however, were similar between the two groups, indicating that the altered pituitary content of GnRH in captive fish may be a result of altered release from the hypothalamus, rather than deficient synthesis (Steven, 2000; Steven et al., 2000; Mylonas et al., 2010).

Hormonal therapies and maturation induction

There are several methods available for spawning induction:

Injection of pituitary extracts, human chorionic gonadotropin (HCG), gonadotropin (GTH), gonadotropin-releasing hormone (GnRH) and GnRH agonists, luteinizing hormone releasing hormone (LHRH) and LHRH agonists (LHRHa).

GnRH and its agonists for spawning induction therapies have important advantages over the use of GTH preparations. GnRH can provide a more balanced stimulation of reproduction events and, better integration of the events with other physiological functions (Lam, 1982; Yaron, 1995; Mylonas et al., 2010). Supplying fresh extracts of the pituitary gland are expensive, difficult availability, variability of the product quality and to obtain the hypophyses the fish should be killed. Also there is possible interference of the hormone administered with the endocrine path of the animal. Besides, they were active only on fish phylogenetically close to the donors. For this reason, GtHs were replaced with human chorionic gonadotropin (hCG), that were characterised by a wide availability on the market and a higher chemical purity, ensuring a better efficacy. In the other hand, luteinizing hormone-releasing hormone analogue des-Gly10(D-Ala6)LHRH- Ethylamide (LHRHa) has been successfully used to induce final maturation and synchronize ovulation of many commercially cultured fish (Donaldson and Hunter, 1983; Park et al., 1997; Mylonas et al., 2010).

Administration methods of GnRHa to broodstocks include injection of GnRH or GnRHa sustained release preparations (Crim and Bettles, 1997; Peter and Yu, 1997; Zohar and Mylonas, 2001). It is the most commonly used due to wider availability, higher purity and standardized activity. Also, piscine pituitary extracts and purified LH obtained through chromatographic separation are used (Lam, 1982; Donaldson and Hunter, 1983; Zohar, 1989b). It has been tested with variable success in many commercially important fish, often in combination with GnRHa. It has been suggested that due to the large molecular size and heterologous nature of GtH preparations, fish develop an immune response and attempts to use it for more than one reproductive season

either fail or require an increasingly higher dose (Zohar, 1989b; Peter and Yu, 1997; Zohar and Mylonas, 2001). GnRH analogues are used in a group of fish species successfully. In carp by the use of GnRH analogues, together with antidopaminergic drugs (pimozide, domperidone, etc.), was successful. Antidopaminergic drugs (are well-known neuroleptic compounds with an extremely long half-life, exhibiting a wide range of physiological effects. The presence of dopaminergic inhibition has been decisively demonstrated in many cyprinid fish and the African catfish (*Clarias gariepinus*, Clariidae), but appears to be absent or weak in most commercially important marine fish (Zohar et al., 1995b). As a result, spawning induction protocols using a combined GnRHa and DA antagonist treatment are used extensively only for cyprinid fish (Peter et al., 1993; Yaron, 1995; Zohar and Mylonas, 2001).

The action of GnRHa takes place at a higher level of the animal endocrine pathway, stimulating the reproductive events in a more balanced and integrated way, with the other physiological functions correlated to the reproduction cycle (Mylonas et al., 1996). Therefore, GnRH seems stimulates also the release of other pituitary hormones involved in the reproductive cycle, like growth hormone (GH), thyroid stimulating hormone (TSH) and Somatolactine (Mylonas et al., 1998b; Zohar and Mylonas, 2001). In the use of gonadotropin hormone, at the beginning, the effective doses of GnRH or its analogues used to induce ovulation in cultured fish ranged from 1 to 15 mg/kg bw (Donaldson and Hunter, 1983; Zohar and Mylonas, 2001). In the majority of scientific reports concerning fish reproduction published in the 1980s, the doses of GnRH analogues ranged from 50 to 100 µg/kg bw. Presently, recommended doses of GnRH analogues for aquaculture vary between 10 and 50 µg/kg bw depending on the fish species, analogue, and mode of its administration (Yu et al., 1997; Zohar and Mylonas, 2001; Zohar et al., 2010). However, injection of GnRHa does not always result in 100% ovulation (Zohar et al., 1995b). In most reports concerning the effect of GnRH analogues, it was shown that GnRH alone is unable to induce ovulation or that the percentage of ovulated females is extremely low at 10 to 20% (Weil et al., 1980; Kouril et al., 1983; Sokolowska et al., 1984; Podhorec and Kouril, 2009). Due to the strong dopaminergic inhibition of LH secretion, typical for the family Cyprinidae, a majority of trials with ovulation induction using only GnRHa failed (Weil et al., 1980; Sokolowska et al., 1984).

According to the results of other authors (Peter et al., 1988; Yaron, 1995; Heyrati et al., 2007; Hamackova et al., 2001; Kouril et al., 1986) the only exceptions to the strong dopaminergic activity in Cyprinidae are tench (*Tinca tinca*), and rudd (*Scardinius erythrophthalmus*), in which even a dose of 1 µg/kg mGnRHa was able to stimulate ovulation in a small number of females. As a consequence of the identification of DA's role in LH

inhibition in Cyprinidae, Peter et al. (1988) developed the so-called LinPe method using the simultaneous administration of GnRH α and effective dopamine D2receptor antagonist. DI disinhibits dopaminergic effect and strengthens the gonadotropin cell stimulation critical for induction of the reovulatory surge of LH (Podhorec and Kouril, 2009).

In studies of the effect of dopamine in fish, the goldfish were the first to demonstrate the effects of blockage of dopamine actions on GtH-II release and ovulation by determining the response to (D-Ala 6 , Pro 9 -Net)-mammalian GnRH (mGnRH-A) when given on combination with the dopamine receptor antagonist, pimoziide. The presence of dopamine inhibition of GtH-II release has been extended to a number of other teleost species, including:

African catfish, coho salmon, Japanese eel, European eel, rainbow and brown trout and tilapia (Lin and Peter, 1996).

Based on the evidence that the failure of cultured fishes to undergo full spermiation and OM is the result of diminished LH release from the pituitary, manipulations of reproductive function have focused first on the use of exogenous LH preparations that act directly at the level of the gonad, and more recently on GnRH α -with or without DA that releases the endogenous LH stores from pituitary. Endogenous LH, in turn, acts at the level of the gonad to induce steroidogenesis and the process of OM and spermiation.

The results indicate that for the induction of a sustained increase in milt production, especially in individuals that are stripped spawned during the reproductive season, a longterm hormonal therapy is necessary, either through the use of multiple injections or controlled release delivery systems. Many different GnRH α delivery systems have been used to enhance spermiation in cultured fishes, such as Atlantic salmon, rainbow trout, coho salmon and Chinook salmon (Mylonas and Zohar, 2001b). In the European seabass, treatment with GnRH α -delivery systems resulted in increased milt production for 28 to 35 days, compared to 7 days only when a single injection of GnRH α was given (Mañanos et al., 2002; Sorbera et al., 1996; Mylonas et al., 2010). Also in the striped bass, GnRH α delivery systems induced increases in milt production for 14 to 20 days (Mylonas et al., 1997, 1998a; Mylonas et al., 2010). To increase the bioavailability and efficiency of GnRH α , several sustained delivery systems have been used successfully in spawning induction of various species of fish; these include cholesterol pellets, cholesterol, cellulose pellets, ethylenevinyl acetate (EVAC), polylactide-glycolide (PLGA) and fatty acid dimer sebasic acid (Fad-Sa) microspheres (Zohar and Mylonas, 2001; Mylonas and Zohar, 2001b). GnRH α implants have also been used in Atlantic halibut to enhance the quality of the

sperm, which is extremely viscous and exhibits very little spermatozoa motility towards the end of the spawning season (Vermeirssen et al., 2003).

The used of different hormone such as CPE, GnRH and LRH in spawning induction therapies in synchronous different fish species are presented in Table 1. Studying the last two decade due to development of aquaculture, the use of hormones to induce OM and ovulation have been increased. Based on the table, most popular hormone for induction is GnRH. Early application of GnRH α was the synchronization of Om and ovulation in salmonids. Treatment with GnRH α is used 1 to 2 week before spawning. A double injection of 10 to 100 μ g/kg B.W. spaced 3 days apart. GnRH α is given in a priming dose (5 to 10%) and a resolving dose (95 to 90%). The same results single or multiple injection of GnRH α has also been used extensively in other fish species such as common carp, sturgeon fishes, and mullet to synchronize ovulation. In sturgeon priming and resolving injection spaced 10 to 24 h apart, and ovulation is accomplished 24 to 50 h afterwards. A double treatment with CPE was used in sturgeon fish farm to induce OM and ovulation (Yousefian et al., 2010a). In propagation of sturgeon first we should be confirmed maturity of fish. Gonad maturity is detected by taking sample, using a plastic cylindrical probe. High degree of migration of germinal vesicle (G.V.) expressed by polarization Index (PI) indicated spawnable females. PI is the ratio of distance between germinal vesicle and the animal pole to the diameter of the oocyte. PI within the range of 5 to 6 was more advanced in maturity than higher or lower in the study of Yousefian et al., 2010a. In order to take place of OM and ovulation, spawner was stimulated by usage of suspension of powdered acetone dried sturgeon pituitaries at a rate of 45 to 50 mg per female of *Acipenser pesicus* (Yousefian et al., 2010a).

A single treatment with CPE has also been reported to be effective (Williot et al., 2005).

In several fin fishes, DA antagonist is also used, administered together with priming dose. In Tench (*Tinca tinca*) a single injection of 20 μ g/kg GnRH α induced ovulation while in common carp the same amount of the hormone with 5 mg/kg of DA analogonis pimoziide induce ovulation. In the gray mullet (*Mugil cephalus*), two injections of 20 μ g/kg GnRH α together with 15 mg/kg of the DA antagonist metoclopramide were very effective in inducing spawning within 24 h (Aizen et al., 2005). Induce oocyte vitelogenesis and ovulation of Gray mullet (*M. cephalus*), are described in detail by Yousefian et al. (2009), by using of mammalian gonadotrophins and synthetic analog of luteinizing hormone releasing hormone (LRH-A2). For successful spawning, a priming injection for females should be when an average egg diameters is 600 μ m or more. A ripe ova have a thin layer cytoplasm covering the yolk and have a single oil droplet. In this case, the egg diameter was 960 μ m and oil globule 360 μ m. Hydration was observed about 10 h after the

injection of effective dose and spawning within 4 to 6 h after beginning of hydration at $24 \pm 1^\circ\text{C}$ and water salinity 33 to 35 ppt. During this time GSI increased from 17 to 28%. The percent of fertilization was up to 80. A combination of hormone in most cases showed better effectiveness comparing to one single hormone. The study was performed in order to try to provoke the gonadotropin wave and ovulation in *Chalcarburnus chacooides* using GnRH analogue, pituitary extract and dopamine antagonist (Yousefian et al., 2008).

Different doses and injection protocols were applied using a combination of LRH-Aa, metoclopramide and carp pituitary extract. *C. chacooides* show wide individual variation and much slower response to different hormonal stimulation. Saline (0.9% NaCl) injected fish were used as a control group and no ovulation occurred in this group. Based on the spawning ratio, fertilization rate and hatching rate, the combination of LRH-Aa 5 $\mu\text{g}/\text{kg}$, carp pituitary extraction 2.7 mg/kg and metoclopramide 2 mg/kg B.W. doses were found to be more efficient than carp pituitary (4mg/kg). No females ovulated in the treatment group receiving metoclopramide at 10 mg/kg and LRH-Aa alone at 100 $\mu\text{g}/\text{kg}$ (Yousefian et al., 2008).

In the chum salmon (*Oncorhynchus keta*), a single injection of 70 $\mu\text{g}/\text{kg}$ GnRH α together with 700 $\mu\text{g}/\text{kg}$ the DA antagonist pimozide were effective in inducing OM and synchronizations ovulation within one week (Park et al., 2007). In rainbow trout (*Oncorhynchus mykiss*) two injections of 25 $\mu\text{g}/\text{kg}$ GnRH α spaced 1 day apart induce synchronizations of ovulation in 100% of the population 11 days after treatment without affecting broodstock survival and egg quality (Vazirzadeh et al., 2008).

Our recent work on brown trout showed, two injections of 20 $\mu\text{g}/\text{kg}$ GnRH α in combination with 10mg/kg DA antagonist domperidone or metoclopramide spaced 4 days apart, have been synchronized ovulation successfully in the Caspian brown trout (*Salmo trutta caspius*) within 1 to 2 weeks after treatment. Another hormone which is widely used is hCG and are preferred over GnRH α by several scientists. The advantage of hCG is that it acts directly at the level of the gonad and does not require the activation of the pituitary gonadotropes or existence of LH stores. Occasionally, GnRH α is not effective or requires a long time to elicit a response. Therefore, hCG may be more appropriate because it acts much faster, via direct gonadal stimulation, in inducing FOM, spermiation and spawning (Hodson and Sullivan, 1993; Zohar and Mylonas, 2001).

A single injection of hCG at 1000 or 2000 IU/kg was effective in inducing OM and ovulation in the spotted sea bass (*Lateolabrax maculatus*) (Lee and Yang, 2002) and in Japanese catfish (*Silurus asotus*), a single injection of 10,000 IU/kg hCG induced OM and ovulation (Kumakura et al., 2003), in pike perch (*Sander Luciperca*) single or multiple injections of 200 IU/kg hCG (Zakes and Szczepkowski, 2004), and in catfish (*Pseudoplatystoma fasciatum*), treatment with CPE and hCG were both

effective in inducing OM and ovulation (Leonardo et al., 2004). In fishes which migrate to the river for spawning, actually are at final stage of naturatiom, therefore no necessary any hormonal treatment. An example are *Caspoan Kutum* (*Rutilus frisii kutum*) which will migrate to the river and are artificially spawn just by gentile forcing in the belly of male and female (Yousefian and Mousavi, 2008).

CONCLUSION

The number of fish species currently under domestication efforts is rising up, due to the development of commercial aquaculture. For domestication and establish a sustainable aquaculture it is necessary to stimulate the fish and control reproduction processes of fish in captivity to acquire high quality seed. Different species of fish have their own physiological and characteristics in reproduction. Therefore it is suggested for different fish using separate protocol of hormones and for new species to have pre-experiment to illustrate the exact dosage and hormonal combination.

REFERENCE

- Abraham M (1988). Recent trends in research on induced spawning of fish in aquaculture. J. Appl. Ichtyol., 4: 49-64.
- Aizen J, Meiri I, Tzchori I, Levavi-Sivan B, Rosenfeld H (2005). Enhancing spawning in the grey mullet (*Mugil cephalus*) by removal of dopaminergic inhibition. Gen. Comp. Endocrinol., 142: 212-221.
- Billard R, Alagarswami K, Peter R.E, Breton B (1983). Potentialisation par le pimozide des effets du LHRH-A sur la se'cre'tion gonadotrope hypophysaire, l'ovulation et la spermiation chez la carpe commune (*Cyprinus carpio*). C.R. Acad. Sci. (Paris), 296: 181-184.
- Crim LW, Bettles S (1997). Use of GnRH analogues in fish culture. In: Fingerman, M., Nagabhushanam, R. and Thompson MF. (eds.), Recent Advances in Marine Biotechnology: Endocrinology and Reproduction. Oxford & IBH Publishing Co., New Delhi, 1: 369-382.
- Crim LW, Glebe BD (1984). Advancement and synchrony of ovulation in Atlantic salmon with pelleted LHRH analog. Aquaculture, 43: 47-56.
- Cardinaud B, Gilbert JM, Liu F, Sugamori KS, Vincent JD, Niznik HB, Vernier P (1998). Evolution and origin of the diversity of dopamine receptors in vertebrates. Adv. Pharmacol., 42: 936-940.
- Davies B, Swanson P, Bromage N (1995). The effects of photoperiod and temperature on serum GtH I and GtH II and the timing of maturation in the female rainbow trout. In: Goetz, F.W., Thomas, P. (Eds.), Reproductive Physiology of Fish V. Proceedings of the Fifth International Symposium, 02 July-08 July 1995. Fish Symposium 95, Austin, USA, pp. 185-186.
- Dickey JT, Swanson P (1995). Development of RNase protection assays for quantification of gonadotropin (GtH I and GtH II). subunit transcript levels in coho salmon. In: Goetz, F.W., Thomas, P. Eds., Reproductive Physiology of Fish V. Proceedings of the Fifth International Symposium, 02 July-08 July 1995. Fish Symposium 95, Austin, USA, p. 28.
- Donaldson EM, Hunter GA (1983). Induced final maturation, ovulation, and spermiation in cultured fish, pp. 351-403. In: W.S. Hoar and D.J. Randall, editors. Fish Physiology, Vol. 9B. Academic Press, New York, NY.
- Hodson R, Sullivan CV (1993). Induced maturation and spawning of domestic and wild striped bass, *Morone saxatilis* (Walbaum), broodstock with implanted GnRH analogue and injected hCG. Aquacult. Fish. Manage., 24: 389-398.

- Heyrati FP, Mostafavi H, Toloei H, Dorafshan S (2007). Induced spawning of kutum, *Rutilus frisii kutum* (Kamenskii, 1901) using (D-Ala₆, Pro⁹-NEt) GnRHa combined with domperidone. *Aquaculture*, 265: 288-293.
- Hamackova J, Kouril J, Barth T, Lepicova A, Kozak P, Lepic P (2001). Induction of ovulation in rudd (*Scardinius erythrophthalmus*) using hormone preparations. Collections Symposium Series, Institute of Organic Chemistry and Biochemistry of the ASCR, Prague, 4: 87-89.
- Johnson JD, Van Goor F, Wong CJ, Goldberg JI, Chang JP (1999). Two endogenous gonadotropin-releasing hormones generate dissimilar Ca(2+) signals in identified goldfish gonadotropes. *Gen. Comp. Endocrinol.*, 116: 178-191.
- Kebabian JW, Calne DB (1979). Multiple receptors for dopamine. *Nature*, 277: 93-96.
- Kouril J, Barth T, Hamackova J, Flegel M (1986). Induced ovulation in tench (*Tinca tinca* L.) by various LH-RH synthetic analogues: Effect of site of administrative and temperature. *Aquaculture*, 54: 37-44.
- Kah O, Lethimonier Ch, Somoza G, Guilguir L, Vail-lant C, Lareyre JJ (2007). GnRH and GnRH receptors in Metazoa: A historical, comparative and evolutive perspective. *Gen. Comp. Endocrinol.*, 153: 346-364.
- Kah O, Dulka JG, Dubourg P, Thibault J, Peter RE (1987). Neuroanatomical substrate for the inhibition of gonadotropin secretion in goldfish: existence of a dopaminergic preoptico-hypophyseal pathway. *Neuroendocrinology*, 45: 451-458.
- Kouril J, Hamackova J, Barth T, Hulova I, Barthova J (1999). Artificial stripping of large mouth buffalo (*Ictiobus cyprinellus*) using carp pituitary, GnRH analog and dopaminergic inhibitor isofloxythepin. Biologically Active Peptides VI, Collection Symposium Series, Prague, 3: 68-70.
- Kumakura N, Sakai K, Takashima F (2003). Reproductive cycle and human chorionic gonadotropin-induced ovulation in hatchery reared Japanese catfish *Silurus asotus*. *Fish. Sci.*, 69: 495-504.
- Linard B, Anglade I, Bennani S, Salbert G, Navas JM, Bailhache T, Pakdel F, Je'go P, Valotaire Y, Saligaut C, Kah O (1995). Some insights into sex steroid feedback mechanisms in the trout. In: Goetz, F.W., Thomas, P. (Eds.), *Proceedings of the Fifth International Symposium on Reproductive Physiology of Fish*, University of Texas, Austin, TX, USA, pp. 49-51.
- Linard B, Anglade A, Corio M, Navas JM, Pakdel F, Saligaut C, Kah O (1996). Estrogen receptors are expressed in a subset of tyrosine hydroxylase-positive neurons of the anterior preoptic region in the rainbow trout. *Neuroendocrinology*, 63: 156-165.
- Lam TJ (1982). Applications of endocrinology to fish culture. *Can. J. Fish. Aquat. Sci.*, 39: 111-1137.
- Lin HR, Peter RE (1996). Hormones and Spawning in fish. *Asian Fisheries Science*. Manila, Philippines, pp. 21-33
- Lin HR, Kraak GVD, Liang J-Y, Peng C, Li G-Y, Lu L-Z, Zhou X-J, Chang ML, Peter RE (1986). The effects of LHRH analogue and drugs which block the effects of dopamine on gonadotropin secretion and ovulation in fish cultured in China. In: Billard, R., Marcel, J. (Eds.), *Aquaculture of Cyprinids*. INRA, Paris, pp. 139-150.
- Lin HR, Zhou XJ, Van Der Kraak G, Peter RE (1991). Effects of gonadotropin-releasing hormone agonists and dopamine antagonists on gonadotropin secretion and ovulation in Chinese loach, *Paramisgurnus dabryanus*. *Aquaculture*, 95: 139-147.
- Lin HR, Kraak GVD, Liang J-Y, Peng C, Li G-Y, Lu L-Z, Zhou X-J, Chang ML, Peter RE (1986). The effects of LHRH analogue and drugs which block the effects of dopamine on gonadotropin secretion and ovulation in fish cultured in China. In: Billard, R., Marcel, J. Eds., *Aquaculture of Cyprinids*. INRA, Paris, pp. 139-150.
- Leonardo AFG, Romagosa E, Borella MI, Batlouni SR (2004). Induced spawning of hatchery-raised Brazilian catfish, *cachara Pseudoplatystoma fasciatum* (Linnaeus, 1766). *Aquaculture*, 240: 451-461.
- Legendre M, Linhart O, Billard R (1996). Spawning and management of gametes, fertilized eggs and embryos in Siluroidei. *Aquat. Living Resour.*, 9: 59-80.
- Lee W-K, Yang S-W (2002). Relationship between ovarian development and serum levels of gonadal steroid hormones, and induction of oocyte maturation and ovulation in the cultured female Korean spotted sea bass *Lateolabrax maculatus* (Jeom-nong-eo). *Aquaculture*, 207: 169-183.
- Malison JA, Held JA (1996). Reproduction and spawning in walleye (*Stizostedion vitreum*). *J. Appl. Ichthyol.*, 12: 153-156.
- Mylonas CC, Zohar Y (2001b). Use of GnRHa-delivery systems for the control of reproduction in fish. *Rev. Fish Biol. Fish.*, 10: 463-491.
- Mylonas CC, Scott AP, Vermeirssen ELM, Zohar Y (1997). Changes in plasma gonadotropin II and sex steroid hormones, and sperm production of striped bass after treatment with controlled-release gonadotropin-releasing hormone agonist-delivery systems. *Biol. Reprod.*, 57: 669-675.
- Mylonas CC, Zohar Y (2001a). Endocrine regulation and artificial induction of oocyte maturation and spermiation in basses of the genus *Morone*. *Aquaculture*, 202: 205-220.
- Mylonas CC, Magnus Y, Gissis A, Klebanov Y, Zohar Y (1996). Application of controlled-release, GnRHa-delivery systems in commercial production of white bass striped bass hybrids (*sunshine bass*), using captive broodstocks. *Aquaculture*, 140: 265-280.
- Mylonas C.C, Fostier A, Zanuy S (2010). Broodstock management and hormonal manipulations of fish reproduction. *Gen. Comp. Endocrinol.*, 165: 516-534.
- Mylonas CC, Woods LC, Thomas P, Zohar Y (1998b). Endocrine profiles of female striped bass (*Morone saxatilis*) during post-vitellogenesis, and induction of final oocyte maturation via controlled-release GnRHa-delivery systems. *Gen. Comp. Endocrinol.*, 110: 276-289.
- Mylonas CC, Hinshaw JM, Sullivan CV (1992). GnRHa-induced ovulation of brown trout (*Salmo trutta*) and its effects on egg quality. *Aquaculture*, 106: 379-392.
- Mylonas CC Hinshaw J, Sullivan CV (1991). Induced ovulation of brown trout (*Salmo trutta*). Using GnRHa and T3, and hormone treatment effects on progeny development. In: Scott, A.P., Sumpter, J.P., Kime, D.E., Rolfe, M.S. (Eds.), *Proceedings of the Fourth International Symposium on Reproductive Physiology of Fish*. Fish Symposium 91, Sheffield, p. 285.
- Mousavi SE, Yousefian M, Hazaee K (2008). Induction and synchronization of ovulation in Caspian brown trout, *Salmo trutta caspius*, using GnRHa in combination with dopamine antagonist.
- Nieuwenhuys R, Donkelaar HJt, Nicholson C (1998). In: *The Central Nervous System of Vertebrates*. Springer, Berlin, New York.
- Prat F, Sumpter JP, Tyler RC (1996). Validation of radioimmunoassays for two salmon gonadotropins (GtH I and GtH II). and their plasma concentrations throughout the reproductive cycle in male and female rainbow trout (*Oncorhynchus mykiss*). *Biol. Reprod.*, 54: 1375-1382.
- Peter RE, Lin H-R, Van der Kraak GV (1988). Induced ovulation and spawning of cultured freshwater fish in China: Advances in application of GnRH analogues and dopamine antagonists. *Aquaculture*, 74: 1-10.
- Peter RE, Yu KL (1997). Neuroendocrine regulation of ovulation in fish: Basic and applied aspects. *Rev. Fish Biol. Fish.*, 7: 173-197.
- Peter RE, Paulencu CR (1980). Involvement of the preoptic region in gonadotropin release-inhibition in goldfish, *Carassius auratus*. *Neuroendocrinology*, 31: 133-141.
- Peter RE, Lin HR, van der Kraak G, Little M (1993). Releasing hormones, dopamine antagonists and induced spawning. In: Muir, J.F. and Roberts, R.J. (eds.), *Recent Advances in Aquaculture*. Blackwell Scientific Publications, Oxford, 4: 25-30.
- Peter RE, Lin HR, Van Der Kraak G (1988). Induced ovulation and spawning of cultured freshwater fish in China: advances in application of GnRH analogues and Dopamine antagonists. *Aquaculture*, 74: 1-10.
- Planas JV, Swanson P, Dickhoff WW (1991). Regulation of testicular steroidogenesis by coho salmon gonadotropins, GTH I and GTH II, in vitro. In: Scott, A.P., Sumpter, J.P., Kime, D.E., Rolfe, M.S (Eds.), *Proceedings of the Fourth International Symposium on Reproductive Physiology of Fish*, University of East Anglia, Norwich, UK, p. 103.
- Podhorec P, Kouril J (2009). Induction of final oocyte maturation in cyprinidae fish by hypothalamic factors: A review. *Vet. Med.*, 54 (3): 97-110.
- Park WD, Lee CH, Lee CS, Kim DJ, Kim JH, Tamaru CS, Sohn YC (2007). Effects of gonadotropin-releasing hormone analog combined with pimoziide on plasma sex steroid hormones, ovulation and egg quality in fresh water-exposed female chum salmon (*Oncorhynchus*

- keta). *Aquaculture*, 271: 488-497.
- Rainis S, Ballestrazzi R (2005). The control of reproduction in finfish species through GnRH treatments. *Ital. J. Anim. sci.*, 4: 345-353.
- Rebers FE, Hassing GA, Zandbergen MA, Goos HJ, Schulz RW (2000). Regulation of steady-state luteinizing hormone messenger ribonucleic acid levels, de novo synthesis, and release by sex steroids in primary pituitary cell cultures of male African catfish, *Clarias gariepinus*. *Biol. Reprod.*, 62: 864-872.
- Sorbera LA, Mylonas CC, Zanuy S, Carillo M, Zohar Y (1996). Sustained administration of GnRH_a increases milt volume without altering sperm counts in the sea bass. *J. Exp. Zool.*, 276: 361-368.
- Sokolowska M, Peter RE, Nahorniak CS, Nahorniak CH, Chang JP (1984). Induction of ovulation in goldfish, *Carassius auratus*, by pimozone and analogues of LH-RH. *Aquaculture*, 36: 71-83.
- Sukumasavin N, Sakulthong S, Sangthong R (2000). A comparison of the potency of dopamine antagonists on spawning induction in Thai carp (*Puntius goni-onotus Bleeker*). *Kasetsart J.*, 34: 240-247.
- Slater CH, Schreck CB, Amend DF (1995). GnRH_a injection accelerates final maturation and ovulation/spermiation of sockeye salmon, *Oncorhynchus nerka*, in both fresh and salt water. *Aquaculture*, 130: 279-285.
- Sokolowska M, Peter RE, Nahorniak CS, Pan CH, Chang JP, Crim LW, Weil C (1984). Induction of ovulation in goldfish, *Carassius auratus*, by pimozone and analogues of LH-RH. *Aquaculture*, 36: 71-83.
- Sullivan CV, Bernard MG, Hara A, Dickhoff WW (1989). Thyroid hormones in trout reproduction: enhancement of GnRH_a and partially purified salmon GtH-induced ovarian maturation *in vivo* and *in vitro*. *J. Exp. Zool.*, 250: 188-195.
- Saligaut et al (1999). Brain aminergic systems in salmonids and other teleosts in relation to steroid feedback and gonadotropin release. *Aquaculture*, 177: 13-20.
- Sakai N, Ueda H, Suzuki N, Nagahama Y (1989). Steroid production by Amago salmon (*Oncorhynchus rhodurus*). Testes at different developmental stages. *Gen. Comp. Endocrinol.*, 75: 231-240.
- Steven C (2000). Studies on the GnRH-GtH system of female striped bass (*Morone saxatilis*): effects of GnRH agonist therapy and comparison of reproductive endocrine parameters between wild and captive fish. Marine Estuarine and Environmental Sciences. University of Maryland, College Park.
- Steven C, Gothlif Y, Holland MCH, Stubblefield J, Mylonas CC, Zohar Y (2000). Differential expression of the three GnRH genes in wild and captive striped bass, *Morone saxatilis*, in response to natural and hormonally induced maturation. In: Norberg, B., Kjesbu, O.S., Taranger, G.L., Andersson, E., Stefansson, S.O. (Eds.), *Reproductive Physiology of Fish*. University of Bergen, Bergen, p. 66.
- Senthilkumaran B, Joy KP (1995). Changes in hypothalamic catecholamines, dopamine- β -hydroxylase, and phenylethanolamine-N-methyltransferase in the catfish *Heteropneustes fossilis* in relation to season, raised photoperiod and temperature, ovariectomy, and estradiol-17- β replacement. *Gen. Comp. Endocrinol.*, 97: 121-134.
- Schulz RW, Andriske M, Lembke PJ, Blüm V (1991). Sex steroids in rainbow trout: plasma levels and testicular secretion *in vitro*. In: Scott, A.P., Sumpter, J.P., Kime, D.E., Rolfe, M.S. (Eds.), *Proceedings of the Fourth International Symposium on Reproductive Physiology of Fish*, University of East Anglia, Norwich, UK, pp. 83-85.
- Schulz RW, Andriske M, Lembke PJ, Blüm V (1992). Effect of salmon gonadotropic hormone on sex steroids in male rainbow trout: plasma levels and testicular secretion *in vitro*. *J. Comp. Physiol. B*, 162: 224-230.
- Vazirzadeh A, Hajimoradloo AM, Esmaeili HR, Akhlaghi M (2008). Effects of emulsified versus saline administration of GnRH_a on induction of ovulation in rainbow trout, *Oncorhynchus mykiss*. *Aquaculture*, 280: 267-269.
- Vermeirssen ELM, Mazorra de Quero C, Shields RJ, Norberg B, Kime DE, Scott AP (2003). Fertility and motility of sperm from Atlantic halibut (*Hippoglossus hippoglossus*) in relation to dose and timing of gonadotropin-releasing hormone agonist implant. *Aquaculture*, 230: 547-567.
- Van der Kraak G, Donaldson EN, Chang JP (1986). Dopamine involvement in the regulation of gonadotropin secretion in coho salmon. *Can. J. Zool.*, 64: 1245-1248.
- Williot P, Brun R, Rouault T, Pelard M, Mercier D, Ludwig A (2005). Artificial spawning in cultured sterlet sturgeon, *Acipenser ruthenus* L., with special emphasis on hermaphrodites. *Aquaculture*, 246: 263-273.
- Weil C, Fostier A, Horvath L, Marlot S, Berscenyi M (1980). Profiles of plasma gonadotropin and 17- β -estradiol in the common carp, *Cyprinus carpio* L., as related to spawning induced by hypophyztation or LH-RH treatment. *Reprod. Nutr. Dev.*, 20: 1041-1050.
- Yaron Z (1995). Endocrine control of gametogenesis and spawning induction in the carp. *Aquaculture*, 129: 49-73.
- Yaron Z, Sivan B (2006). Reproduction, pp. 343-386. In: Evans D.H., Claibourne J.B. (eds.): *The Physiology of Fishes*. CRC Press, Boca Raton, p. 601.
- Yousefian M, Ghanei M, Pourgolam R, Rostami HKh (2009). Gonad Development and Hormonal Induction in Artificial Propagation of Grey Mullet, *Mugil Cephalus* L. *Res. J. Fish. Hydrobiol.*, 4(2): 35-40.
- Yousefian M, Gezel HG, Hedayatifard M (2008). Induction of ovulation in endemic *Chalcarbunus chalcoides*, living in the Caspian Sea, using LRH-Aa combined with metoclopramide. *Afr. J. Biotechnol.*, 7(22): 4199-4201.
- Yousefian M, Mosavi H (2008). Spawning of South Caspian Kutum (*Rutilus frisii kutum*) in Most Migratory River of South Caspian Sea. *Asian J. Anim. Vet. Adv.*
- Yousefian M, Najafpour Sh, Farabi SV, Najafpour GD (2010a). Artificial Spawning and Early Development of *Acipenser persicus*. *World J. Fish Mar. Sci.*, 2(3): 258-263.
- Yousefian M (2010b). Stock Identification and Genetic Variation at Microsatellite Loci of Caspian Sea Salmon (*Salmo trutta caspius*). *World J. Fish Mar. Sci.*, 2(6): 508-512.
- Yu KL, Lin XW, da Cunha Bastos J, Peter RE (1997). Neural regulation of GnRH in teleost fishes. In: Parhar, I.S., Sakuma, Y. (Eds.), *GnRH Neurons: Gene to Behavior*. Brain Shuppan, Tokyo, pp. 277-312.
- Zakes Z, Szczepkowski M (2004). Induction of out-of-season spawning of pikeperch, *Sander lucioperca* (L.). *Aquacult. Int.*, 12: 11-18.
- Zohar Y, Muñoz-Cueto JA, Elizur A, Kah O (2010). Neuroendocrinology of reproduction in teleost fish. *Gen. Comp. Endocrinol.*, 165: 438-455.
- Zohar Y (1998a). Hormone profiles of captive striped bass (*Morone saxatilis*), during spermiation, and long-term enhancement of milt production. *J. World Aquacult. Soc.*, 29: 379-392.
- Zohar Y, Tosky M, Pagelson G, Finkelman Y (1989b). Induction of spawning in the gilthead seabream, *Sparus aurata*, using wD-Ala6-Pro9NETx-LHRH: comparison with the use of HCG. *Israeli J. Aquacult-Bamidgeh*, 41: 105-113.
- Zohar Y, Mylonas CC (2001). Endocrine manipulations of spawning in cultured fish: From hormones to genes. *Aquaculture*, 197: 99-136.
- Zohar Y, Harel M, Hassin S, Tandler A (1995b) Gilthead sea bream (*Sparus aurata*). In: Bromage, N.R. and Roberts, R.J.(eds.), *Broodstock Management and Egg and Larval Quality*. Blackwell Science, Oxford, pp. 94-117.
- Zohar Y, Elizur A, Sherwood NM, Powell JF, Rivier JE, Zmora N (1995a). Gonadotropin-releasing activities of the three native forms of gonadotropin-releasing hormone present in the brain of gilthead seabream, *Sparus aurata*. *Gen. Comp. Endocrinol.*, 97: 289-299.
- Zohar Y (1989a). Endocrinology and fish farming: Aspects in reproduction growth, and smoltification. *Fish Physiol. Biochem.*, 7: 395-405.
- Zohar Y (1996). New approaches for the manipulation of ovulation and spawning in farmed fish. *Bull. Natl. Res. Inst. Aquacult. Suppl.*, 2: 43-48.