

Review

Algae as Antioxidants and Effective Fish Feed a Review

Priyanka Mukherjee*, Ruma Pal

Department of Botany, Phycology laboratory, University of
Calcutta, India.

Accepted 15 December, 2020

The oxidative stress in the body of an organism whether plant or animal leads to the production of several reactive oxygen and reactive nitrogen species abbreviated as ROS and RNS respectively. Algae, because of the presence of various biochemical compounds can act as a potent antioxidative agent and can thrive against the stressful conditions produced due to the oxidative stress. These secondary metabolites are high value compounds which can be efficiently used in the field of aquaculture as feed for different fishes. Thus in this review we have discussed the various compounds isolated from algae which allows the algal life to thrive in high oxidative stress as well as we have discussed different potential algal genera being used as feed for different fishes.

Key words: Reproduction induced spawning, dopamine antagonists, GTH-II- HCG.

INTRODUCTION

The imbalance between the pro-oxidants and the antioxidants leads to the production of several reactive oxygen species commonly called as ROS. These reactive oxygen species are mainly hydroxyl group (OH⁻), superoxide anion (O₂⁻), hydrogen peroxide (H₂O₂) and proxy radicals. These reactive oxygen species in low concentration generally bring about minor changes in the normal cellular redox status, immune function and intracellular functioning of a particular organism. But high concentration of reactive oxygen species leads to the peroxidation of lipids, damages protein and DNA and gets involved with the cellular function leading to cell death by necrosis and cellular apoptosis. Along with damage of the cellular components, ROS also plays an active role in regulation of signal transduction system, gene expression, activation of receptors and nuclear transcription factors. Algae are exposed to a combination of diverse environmental conditions which often may act as stress and lead to the production of different free radicals. Thereby, their elements of the photosynthetic apparatus are also being exposed to photodynamic damage regularly, because the major constituent of the thylakoid membrane is polyunsaturated fatty acids. But the absence of damage to algae in spite of being in proximity to the ROS suggests that these cells have protective antioxidant mechanisms and compounds which provide them shelter from harmful activity of ROS [1].

The protective compounds present in algae are majorly polysaccharides, fattyacids, tannins, pigments, proteins and vitamins. The phlorotannins (compounds with polymerized phloroglucinol units) are also noteworthy antioxidants present in algae especially in Phaeophyceae. They find profound usage in broad therapeutic perspectives, such as anti-diabetes, antioxidation, radiation protection, anti-cancer, anti-HIV as well as anti-allergic activities. Other than polyphenols algae are also rich in peptides which are gaining interest because of their potent antioxidative nature. Thus these algal peptides can be easily implemented as alternative antioxidative supplements [2]. Algae are also rich in phytophenolic compounds and pigments. The Rhodophyta (red algae) are a distinct eukaryotic lineage, which are rich in accessory photosynthetic pigments such as phycoerythrin, phycocyanin and allophycocyanins arranged in phycobilisomes. They are also efficient antioxidants. Thus as algae, have been documented to be rich sources of various bioactive compounds they can easily and safely be incorporated in various biomedical fields and food industry. In animal system the ROS bring about oxidative modification of DNA and proteins and other cellular molecules leading to the development of various ranges of common diseases such as cardiovascular disease, atherosclerosis, diabetes mellitus inflammatory conditions, and neurodegenerative disease such as Alzheimer's disease, Parkinson's disease and immune dysfunction.

*Corresponding author. E-mail: priyankamukherjee76@gmail.com.

Mutations and cancer and age related degenerative disease. Thus to stop the damage causing activity of ROS an efficient antioxidant is required. The use of synthetic antioxidants has decreased nowadays because studies have shown that synthetic antioxidants act as promoters of carcinogenesis furthermore people today reject to use synthetic food additives frequently. Thus these findings along with the general interest have promoted the need for finding out new alternative antioxidants from natural sources. Sea weeds have found their usage since many years in the folk medicine for variety of remedial purposes [3].

Microalgae also play an important role as feed ingredients for fishes, because of their nutritive properties, vibrant pigment profiles and a high content of bioactive compounds with cytostatic, antiviral, anthelmintic, antifungal and antibacterial activities as detected in green, brown and red algae. Aquaculture or rearing of aquatic organisms under controlled conditions for commercial purposes date back to 6000 BC. The concept was first developed by Gunditjmarra people of Victoria, Australia where they first converted volcanic floodplains in the vicinity of Lake Condah into complex of channels and dams for capturing eel fish for consumption. Today, almost 50% of the world's trade comes from the aquaculture based industry. Fish culture in India is very popular and extensive. Along with the countries consumption, these fishes also enter the world market as a form of export. Intensive aquaculture is highly dependent on fish oil and fish meal as feed ingredients. Fish meal stands to be essential dietary sources of protein, amino acids, and fatty acids. To avoid different constrains in this field the usage of algae as a potent feed has found different openings. Several studies have revealed that different algae are being commonly used in the field of aquaculture as different fish feed. are *Chlorella*, *Dunaliella*, *Haematococcus*, *Ulva*, *Enteromorpha*, *Laminaria*, *Porphyra*, *Sargassum*, *Gracillaria* and many more [4].

This review deals with the different antioxidant contents of different algae and the use of different algae with high antioxidant properties in the field of aquaculture.

Compounds responsible for antioxidative property in algae

In recent years, functional foods have been noticed to gain more interest because of their ability to provide physiological benefits in addition to nutritional and energetic, as, for instance, antihypertensive, antioxidant or anti-inflammatory actions. A functional food can be defined as a food that benefits in one or more physiological functions, increases the welfare and/or decreases the risk of suffering from a particular disease. Along with the functional food the most upcoming field in the recent years are the nutraceutical aspects of natural products. People today are more interested or rather prefer natural products both as functional food as well as nutraceuticals. Today the natural sources which are being widely exploited for these purposes are plants, food by-products, fungi and algae. Algae nowadays have become a hot topic of discussion because of their high content of biochemical parameters such as polyunsaturated fatty acids, β -carotene and other pigments (antioxidants), sulphated polysaccharides (antiviral), and sterols (antimicrobials) [5].

Several studies have shown that algae have developed an efficient defense system that helps them to survive under environmentally adverse conditions during evolution. This property has developed in algae due to the compounds present such as tannins, peptides, pheophytins polysaccharides and other pigments.

Tannins are phenolic compounds that show antioxidant activity due to their high redox properties, which play an important role in absorbing and neutralizing free radicals thereby quenching singlet and triplet oxygen or decomposing peroxides. Studies shown that enzymatic hydrolysate from brown alga, *Ecklonia cava* collected from Jeju Island coast of Korea to have a high antiproliferative and antioxidative action. This algal enzymatic hydrolysates had efficient antioxidative action against the DPPH reagent as well as superoxide anions, hydrogen peroxides, hydroxyl radicals which act as the precursors of the singlet oxygen species, thereby indirectly inhibiting lipid peroxidation. This property was attributed to the algae by the various polyphenolic as well as polysaccharide compounds present in it. The crude polysaccharide and crude polyphenolic fractions of the alga showed antioxidative action. Different studies have shown that in *E. cava*. it is the polyphenolic compounds including phlorotannins which remains to be majorly responsible for thriving against various oxidative stress and damages in algae.

Polyphenols have also been found to be widely present in different algae such as *Sargassum swartzii*, *Cystoseira myrica*, *Colpomenia sinuosa*. Algal polyphenolic compounds apart from acting as a good antioxidant in various cell lines also show effective activity in delaying fish oil rancidity. This was first shown by Yan in 1996, where Phlorotannin isolated from *Sargassum kjellmanianum* could also protect fish oil from rancidity. This is due to the fact that polyphenols easily transfer a hydrogen atom to lipid peroxy cycle and form the aryloxy, which being incapable of acting as a chain carrier, couples with another radical thus quenching the radical process [6].

Apart from the polyphenols the pigments also play a major role in antioxidative property of algae. Microalgae are rich sources of pigments comprising of mainly two main types of chlorophyll, chlorophyll a and chlorophyll, β -carotene, astaxanthin and lutein which are considered to be most important forms of 600 naturally occurring carotenoid for their potential applications in food additives for human nutrition, nutraceuticals, nutritional supplements. Carotenoids are the only group of biologically active compounds, that possess antibacterial, antiviral, antifungal, antioxidative, anti-inflammatory, and antitumor properties. Several algal species such as *Haematococcus pluvialis*, *Dunaliella salina*, *Chlorella* sps, *Scenedesmus* sps, *Spirulina platensis*, *Botryococcus braunii*, and Diatoms have gained world wide fame in the commercial production of β -carotene, lutein, canthaxanthin, astaxanthin, and fucoxanthin [7].

The algal pigments such as astaxanthin also plays a very important role as free radical scavenger in algae such as *Haematococcus pluvialis*. Research work has shown that the same mechanism also occurs in *Dunaliella bardawil*. Carotenoids such as astaxanthin act as scavenger of several active oxygen species such as free oxygen radical, hydrogen peroxide, peroxy radicals and hydroxyl radicals ($\text{HO}\cdot$) both in vitro and in vivo. Thus, astaxanthin, provides protection to the cytoplasm and lipid membranes. Therefore, this particular carotenoid is now being utilized as a pigment source in pisciculture and is also being investigated for medical applications because it is more antioxidative than β -carotene and vitamin E.

Other than the pigments algae are also rich in polysaccharide also show antioxidative activity. Different studies of Maximum works with brown algae has been reported till date but only little information is available with Chlorophycean members.

ALGAE AS FISH FEED ADDITIVES

Aqua farming today is also one of the most important businesses all over the world. Use of algal feed in fish aquaculture is well in practice for both edible and colored fishes for their nutritional constituents as well as pigment composition. Works of several authors have shown the usage of different algae such as Porphyridium, Pavlova, Chaetoceros, Gracillaria, Palmaria as efficient color elicitors and growth enhancers in cichlid fish, rainbow trout, fish larvae, bivalve mollusks, black tiger prawns and several gastropods. Several genera like Skeletonema, Chaetoceros, Thalassiosira, Tetraselmis and Isochrysis with long-chain polyunsaturated fatty acids are also well documented as fish feed ingredients or live feed.

Studies have shown use of different algal genera, such as Isochrysis, Pavlova, Chaetoceros, Gracillaria, Palmaria, Haematococcus, Arthrospira, Dunaliella, Chlorella, Chlorococcum etc. to play important role as food ingredients for fishes as they are rich sources of several bioactive compounds which help them. The rich contents of several pigments especially carotenoids have made algae popular amongst fish breeders as feed ingredients. Studies revealed that from the green algae Haematococcus pluvialis, β , β -carotene and lycopene can be extracted and used as an efficient alternative to synthetic pigments in red porgy.

A green algae Ulva containing good vitamin, mineral and pigment profiles have found its usage as a potent feed ingredient for a wide range of fish species. Incorporation of Cladophora, Chaetomorpha, Pithophora, Rhizoclonium, Spirulina, Leptolyngbya etc in fish feed enhanced the carotenoid content as well as immunity in goldfish as well as catfish respectively. Several studies have shown the usage of different algae such as Leptolyngbya valderiana, L. tenuis, Arthrospira maxima, Navicula minima, Nostoc ellipsosporum, Ulva, Cytoseira, Chlorella, Porphyridium, Isochrysis, Pavlova, Chaetoceros, Gracillaria, Palmaria, Isochrysis etc. as efficient color elicitors in cichlid fish, golden gourami, wag swordtail, orange

molly pink zebra, rainbow trout, tetras, prawns, Nile tilapia etc. Therefore incorporation of varied algae in the fish meal diet is cost effective way is a major challenge for improvement of both the production rate as well as their immunity.

CONCLUSION

Researches through years have thus shown algae to be efficient multitasking organisms. They not only play major role in pollution controlling system but also act as efficient antioxidant against oxidative stress management. The high biochemical contents have always proved algae to be the best natural resource ever to be found. Thus because of these properties algae would find immense usage in medicinal field as well as in the field of aquaculture [8].

REFERENCE

1. Sousa I, Gouveia L, Batista A P, Raymundo A, Bandarra NM (2008). Microalgae in novel food products. *Food chem res devel.* 4: 75-112..
2. Coulombier N, Jauffrais T, Lebouvier N. (2021). Antioxidant Compounds from Microalgae: A Review. *Mar drug.* 19 : 549.
3. Hudek K., Davis LC, Ibbini J, Erickson L (2014). Commercial products from algae. *Alg Bioref.* 5: 275-295.
4. Rizwan M, Mujtaba G, Memon SA, Lee K, Rashid N (2018). Exploring the potential of microalgae for new biotechnology applications and beyond: a review. *Renew Sustain Ener Rev*, 92, 394-404.
5. Yaakob Z, Ali E, Zainal A, Mohamad M, Takriff MS. (2014). An overview: biomolecules from microalgae for animal feed and aquaculture. *Journal of Biological Research-Thessaloniki*, 21, 1-10.
6. Ansari FA, Guldhe A, Gupta SK, Rawat I, Bux F (2021). Improving the feasibility of aquaculture feed by using microalgae. *Envir Sci Poll Res*, 28, 43234-43257.
7. Camacho F, Macedo A, Malcata F (2019). Potential industrial applications and commercialization of microalgae in the functional food and feed industries: A short review. *Marine drugs*, 17, 312.
8. Sharma P, Sharma N (2017). Industrial and biotechnological applications of algae: a review. *J Adv Pla Bio*, 1, 01.