

Editorial

Caring of sow and piglets

Federico Agostini*

Department of Medical Genetics, University of Calgary, Canada

Accepted 10 December 2021

EDITORIAL NOTE

A sow ought to have something like fourteen nipples which ought to be long and slim enough for the piglet to get a handle on. Every piglet nurses from its own nipple taking care of consistently. The previously conceived and more grounded piglets utilize the nipples closest the sow's head which produce amount of milk. As a sow gets more seasoned and has more lifters the nipples can turn out to be huge making it hard for the piglet to nurse. In some cases the back nipples don't create a lot of milk. A sow might be not able to take care of all her young and is not, at this point fit for rearing. A sow can experience the ill effects of mastitis which may create because of harm to the nipples brought about by the piglet's teeth. Cutting the teeth of the piglet forestalls slices to the nipples (Ocepek M et al., 2018), (Ocepek M et al., 2017).

Not the entirety of the piglets will develop at a similar rate, some will be conceived more modest than the others. They battle for feed and the more modest piglets will develop at a more slow rate and even bite the dust. You can hope to see a distinction in weight gain and development between the individuals from any litter, however on the off chance that the entirety of the piglets don't develop well and there are no undeniable indications of infection you should presume helpless milk creation by the mother. This is regularly the situation with old sows. It can get important to encourage the piglets, for example put them with an alternate sow for taking care of. It is fundamental for all piglets to take colostrum from the mother. They will take the primary feed inside 1 hour of being conceived. On the off chance that a sow passes on during farrowing her lifter can be cultivated to another. (Kirkden RD et al., 2013), (Gustafsson M, et al., 1999).

A sow may kick the bucket and there could be no non-permanent mother accessible. The litter can be raised by hand taking care of. To hand back a litter the accompanying will be required: Taking care of jugs and nipples (areolas) which are altogether cleaned between each taking care. A spotless dry box containing clean sheet material for the infant piglets which can be kept in a warm spot. Ordinary feeds should be given at time periods to 2 hours (Jarvis S, et al., 1999).

REFERENCE

1. Ocepek M, Andersen IL. (2018). Sow communication with piglets while being active is a good predictor of maternal skills, piglet survival and litter quality in three different breeds of domestic pigs." (*Sus scrofa domestica*). PLoS One. 13(11):e0206128.
2. Ocepek M, Rosvold EM, Andersen-Ranberg I, Andersen IL. (2017). Can we improve maternal care in sows? Maternal behavioral traits important for piglet survival in loose-housed sow herds." *J Animal Sci.* 95(11):4708-4717.
3. Kirkden RD, Broom DM, Andersen IL. (2013). Invited review: Piglet mortality: Management solutions." *J Animal Sci.* 91(7):3361-3389.
4. Gustafsson M, Jensen P, de Jonge FH, Illmann G, Spinka M. (1999). Maternal behaviour of domestic sows and crosses between domestic sows and wild boar." *Appl Animal Behav Sci.* 65(1):29-42.
5. Jarvis S, McLean KA, Calvert SK, Deans LA, Chirnside J, Lawrence AB. (1999). The responsiveness of sows to their piglets in relation to the length of parturition and the involvement of endogenous opioids." *Appl Animal Behav Sci.* 63(3):195-207.

*Corresponding author. Federico Agostini
E-mail: federico.agostini@11scilifelab.se