

African Journal of Infectious Diseases Research ISSN 2756-3340 Vol. 7 (9), pp. 001-006, September, 2020. Available online at www.internationalscholarsjournals.org © International Scholars Journals

Author(s) retain the copyright of this article.

Full Length Research Paper

Comparative biochemical and molecular evaluation of swarming of *Proteus* and effects of anti-swarm agents

Iwalokun BA¹*, Olukosi YA², Adejoro A¹, Olaye JA¹, Fashade O¹

¹Biochemistry Department, Lagos state University, P.M.B. 1087, Apapa, Lagos, Nigeria. ²Genetics Division, Nigerian Institute of Medical Research (NIMR), 6 Edmond Crescent, Yaba – Lagos, P.M.B. 2013, Lagos, Nigeria.

Accepted 16 April, 2020

In addition to inadequate understanding of swarming motility and virulence of *Proteus*, there is paucity of information on the relative effectiveness of the various anti-swarm agents. The anti-swarming effects of urea, sodium dodecylsulphate (SDS) and trihydroxymethylglycine (Tris) on 40 clinical isolates of *Proteus* Spp. were comparatively investigated and plasmids associated with swarming were characterized. The three substances elicited a comparable concentration-dependent anti-swarming property at 0.25 - 1.25% on nutrient agar. Anti-swarm agents displayed heterogeneity in their ability to cause significant decreases in the expression of virulence factors. Swarm motility was further found to be strongly associated with the expression of virulence factors in these strains. Of the *Proteus* strains tested, 32 were found to harbour 1 - 4 plasmids of size ranging from 6.0 - 33.5 kb. Plasmid curing resulted in loss of swarming in 65.6% of these strains. In order to reduce the risk of infection with virulent *Proteus* strains, the laboratory use of urea and SDS is suggested.

Key words: Proteus, swarming, urea, SDS, Tris.

INTRODUCTION

Proteus species contribute greatly to neonatal mortality among Africans with urinary tract infections and are a leading cause of bacteruria associated pregnancy loss worldwide (Orrett, 1999; Cutner et al., 1992). In Nigeria, *Proteus* strains are frequently isolated from patients with kidney diseases, benign prostate hypertrophy and indwelling catheters and have assumed a leading role next to *Pseudomonas aeruginosa* as potential nosocomial pathogens in most Nigerian hospitals and a threat to public health (Kesah et al., 1992; Adeyemo, 1994). Cases of re-current bacteruria are very common in developing countries in particular and this have been blamed on culture diagnostic inappropriateness caused by swarming motility of *Proteus* and related organisms (Williams and Schwartzhoff, 1978; Harshey, 1994). VanAsten and Gaastra (1999) reported difficulties in identifying *P. aeruginosa* and *Streptoccocus G* in mixed cultures involving *Proteus*. This scenario may contribute to spread of multidrug resistance strains and treatment failures of polymicromicrobial infections. Efforts to annul the effects of swarming and enhance diagnosis through culturing in minimal media often fail to identify copathogens with *Proteus* that are fastidious or require extra nutrient supplements for growth (Reynolds and Penn, 1994).

^{*}Corresponding author. E-mail: bamwal@yahoo.com. Phone: 234-1-2601089

However, on solid media many substances have been found to demonstrate anti-swarm properties. They include urea (William, 1973), ethanol and sodium azide (Hernandez and Cavallo, 1999), barbitone and activated charcoal (Alwen and Smith, 1967). Recently, swarming refraining effect of urea was demonstrated on clinical isolates of Proteus from Lagos, and at 0.75 - 1-25%. Urea was found to refrain swarming and allow pneumoniae identification of Klebsiella and Staphyloccocus aureus without compromising extended beta lactamases strain identification (Iwalokun et al., 2003). However, knowledge concerning whether urea influence the expression of virulence factors among these strains is not yet advanced. Elsewhere, multiple drug resistant and biofilm producing swarmed cells have been found to show sensitivity to sodium dodecyl sulphate (SDS) (Davies et al., 1998). The latter seem to support our observation that SDS also refrained swarming greatly in P. aeruginosa and Proteus. Trihydroxymethylglycine (Tris) is widely used in buffer systems involving microbial culture and enzymology, biotechnology and genetic engineering but there have not been clues to show whether this substance could inhibit translocation of bacteria on solid media in vitro. In developing countries, anti-swarm agents are ideally expected to be highly effective, readily available for routine use and confer reduction in the risk of transmission of pathogenic organisms.

Generally, the roles of plasmids in swarming motility have experimentally been demonstrated (Belas et al., 1998) but there is still insufficient information on plasmid profiles in local clinical isolates of *Proteus*. Such information could increase understanding on whether swarming-related genes are plasmid borne.

Hence the present study, which enrolled forty *Proteus* strains, was designed to compare and evaluate the biochemical effects of SDS, urea and Tris on total protein, carbohydrate content, cellular RNA and the expression of extracellular protease in *Proteus*. The involvement of plasmids in swarm motility in these isolates was also determined.

MATERIALS AND METHODS

Proteus strains

A panel of 40 newly isolated *Proteus* strains showing swarming motility on nutrient and broth agar plates were recruited for study. The isolates were obtained from hospital samples (urine, wound exudates, discharging ear specimen, vagina discharge and diarrhoiec stool) and identified in the Microbiology and Genetic Department of the Nigerian Institute for Medical Research, Lagos, Nigeria according to Cowan (1974)

Effects of Tris, urea and SDS on swarming

Cultures were grown to late exponential phase ($A_{625nm} = 0.7 - 0.9$) and 2 µl of 10⁸ cfu/ml of each *Proteus* strain was inoculated at the center of grown in nutrient agar medium containing 0.3% meat extract, 1% peptone and 1.8% Eiken agar and supplemented with Tris, SDS or urea at 0.25 - 1.25% concentrations. Inoculated plates lacking Tris, SDS and urea were also prepared and used as positive controls. The plates were then incubated at 37^{0} C for 24 h under aerobic conditions. The outer diameter of swarming zone from the point of inoculation was measured in millimeter (mm).

Cell- free protein determination

An overnight nutrient broth culture of each of the *Proteus* strains was centrifuged at 2000 x g for 10 min and the resulting supernatant was used for total protein determination according to Lowry et al. (1951). Bovine serum albumin ($500 - 3500 \mu g/ml$) was used as a standard.

Effects of Tris, urea and SDS on Carbohydrate content

The extracellularly secreted carbohydrates in the *Proteus* strains were quantitated from culture supernatants using the anthrone method (Wharton and McCarty, 1972). The secreted carbohydrate was determined in the presence and absence of the anti-swarming agents.

Effects of Tris, urea and SDS on cellular RNA concentration

Cellular RNA content was estimated following cell lysis according to Jayaraman (1988). Cells were lysed by the addition of 400 μ l 4% SDS in Tris EDTA containing 0.4 N NaOH. 300 μ l of sodium acetate buffer (pH 5.2) was added five minutes later to stop lysis. Cell debris was removed by centrifugation at 3000 x g for 10 min. Yeast RNA (150 - 200 μ g/ml) was employed for calibration curve preparation.

Urease assay

One ml aliquot of Proteus culture supernatant was added to 1ml of 0.2 M phosphate buffer containing 3% urea and incubated for 15 min at 55^{0} C. Reactions were terminated by quick transfer to ice and addition of 1 ml 0.66 N H₂SO₄. Protein was estimated according to Lowry et al. (1951) and the liberated ammonia was determined colorimetrically at 500 nm using Nessler's reagent (Jayaraman, 1988). Specific activity of urease was measured as moles of ammonia liberated/mg protein/min.

Protease assay

Extracellular protease in broth culture supernatants of the *Proteus* strains treated and untreated with the anti-swarming agents was assayed for using the protocol of Green et al. (1989). One unit of enzyme activity was defined as the proteolysis of 1 μ g of azocasein substrate per hour at pH 7.0 and 25⁰C.

Plasmid extraction

Plasmid DNAs from *Proteus* strains treated and untreated with antiswarm agents were extracted using the alkaline lysis procedure of Takahashi and Nagano (1976). The extracted plasmids were electrophoresed on 0.8% ethidium bromide pre-stained agarose gel with tris-borate buffer (pH 8.6). Plasmid DNAs were electrophoresed in parallel with standard molecular weight markers (3.5 – 25.2 kb) to allow extrapolation of their sizes.

Plasmid curing

Proteus strains harbouring plasmids were cured of their plasmids according to Garriga et al. (1993). An overnight culture of each *Proteus* strain standardized to 1 x 10^8 cfu/ml with phosphate buffered saline (pH 7.2) was subcultured at 5 x 10^5 cfu/ml into 30 ml nutrient broth (pH 7.6) containing 1.25% ethidium bromide solution. Cultures were incubated for 72 h with continuous shaking (120 rpm) under aerobic condition. Cell pellets were then obtained by centrifugation (5000 rpm, 10 min, 4 0 C). Curing is indicated by loss of plasmids and confirmed by agarose gel electrophoresis described earlier. Swarming motility of cured *Proteus* strains was also investigated in nutrient agar as described previously.

Statistical analysis

Data were expressed as means and means \pm standard deviation. Regression analysis was used to predict associations and evaluate relationships between parameters. Generally, relationships or associations with P < 0.05 were considered significant and strong.

RESULTS

The effects of urea, Tris and SDS on swarm motility, extracellular protease and urease activity, cellular RNA and carbohydrate content among the 40 *Proteus* strains tested are presented in Figures 1 - 5. Swarming motility indicated by outer colony diameter of the strains on agar was observed to decrease significantly (P < 0.05) with increasing concentration of the anti-swarm agents (Figure 1). Concentration dependent reduction in urease activity was also observed with SDS and urea eliciting greater reduction effect on enzyme activity (Figure 2). Similar result was observed in the extracellular protease assay (Figure 3). The anti-swarm agents were also observed to elicit comparable reduction in cellular carbohydrate and RNA levels in the *Proteus* strains tested (Figures 4 and 5).



Figure 1. Swarming inhibitory property of SDS, Tris and urea among *Proteus* isolates from Lagos, Nigeria.



Figure 2. Effects of Tris, urea and SDS on urease activity among the *Proteus* strains.

Figure 3. Effects of SDS, urea and Tris on extracellular protease activity among the *Proteus* strains.

Figure 4. Effects of Tris, urea and SDS on carbohydrate expression among the *Proteus* strains.

Figure 5. Effects of Tris, urea and SDS on cellular RNA level among the *Proteus* strains.

Unlike Tris, SDS and urea generally caused a sharp drop in cellular carbohydrate and RNA as well as extracellular urease and protease activity at 0.5 - 1.0 % concentrations (Figures 2 - 5). Furthermore, regression analysis for the assessment of the relationship between swarm motility inhibition and expression of virulence factors in the tested *Proteus* strains revealed strong association of refrained swarming with protease activity, urease activity, cellular RNA level and carbohydrate expression.

Plasmid analysis revealed that 8 strains were plasmidless, while the remaining strains harboured 1 - 4 plasmids of size ranging from 6.0 - 33.5 kilo base pairs. Treatment with the anti-swarming agents and curing with ethidium bromide were observed to result in plasmid loss (Table 1, Figure 6). Swarm motility was observed in 11 of 32 strains cured of their plasmids.

Table 1.	Plasmid analysis of the Proteus is	solates.

Number of Proteus isolates	No. of plasmids	Plasmid size (Kb)
8	0	-
15	1	6.0, 10.5, 12.5, 14.0
12	2	6.0, 10.5, 12.5, 14.0, 20, 33.5
4	3	6.0, 10.5, 28.0, 33.5
1	4	6.0, 10.5, 33.5

DISCUSSION

Swarming motility, a surface dependent coordinated multicellular behaviour has been established to be the basis for colonization and expression of virulence factors by pathogenic organisms including *Proteus*. This phenotype is a diagnostic constraint and a barrier to adequate anti-proteus measures in humans and animals

Figure 6. Plasmid profile analysis of *Proteus* strains with swarming refrained by Sodium dodecylsulphate, Tris and urea. Lane 1 = Negative control; Lanes 2 and 3 = untreated *Proteus* strains; Lane 4 = *Proteus* strains treated with SDS; Lane 5 = DNA markers (3.5 – 25.2 kb); Lane 6 = Plasmid cured *Proteus* strains; Lane 7 = *Proteus* strains treated with Tris; Lane 8 = *Proteus* strains treated with urea. Band sizes were measured in kilo base pairs (kb).

(Iwalokun et al., 2003; VanAsten and Gaastra, 1999). The present study has provided the molecular and biochemical consequences of the compared swarming refraining properties of urea, SDS and Tris.

The observed swarming refraining action of urea similar to that of SDS is in consonance with previous reports (Williams, 1973; Hernandez and Cavallo, 1999). The ability of SDS to inhibit swarming motility may not be unconnected with the demonstrative evidence of its biocidal effect on signal molecules involved in biofilm formation by P. aeruginosa (Davies et al., 1998). Furthermore, Campanac et al. (2002) recently demonstrated biocide cationic agents as suppressor of biofilm formation in pathogenic organisms. Meanwhile, previous among members of studies the Enterobacteriaceae have successfully demonstrated the coupling of swarming motility to biofilm formation with this relationship strongly established in Yersinia enterocolitica (Young et al., 1999).

To further improve understanding on the consequences of swarming motility inhibition on virulence, we investigated the effect of the anti-swarming agents on urease and protease activity, cellular RNA level and carbohydrate expression in the 40 *Proteus* strains tested. These substances were observed to cause marked reduction in these parameters, contrary to the previously reported roles played by serine, methionine and glutamine (Iwalokun and Akinwunmi, 2002). The reduced cellular RNA may be a manifestation of gene repression mimicking the observation of Rasmussen et al. (2000) that brominated furanones, inhibit swarming genes transcription and the work of Liaw et al. (2003) in which inhibition of virulence gene expression was linked to mRNA degradation in *Proteus mirabilis*. The observed decline in protease and urease activity is supportive to the findings in which p-nitrophenylglycerin was used to inhibit swarming differentiation in *Proteus in vitro* (Liaw et *al*, 2000). Resistant isolates were subsequently found to display super – swarming phenotype (Liaw et al., 2001).

The concentration dependent mode of swarming motility inhibition demonstrated in this study further reveals the possible use of these substances at low concentrations to achieve distinct colony cultures from polymicrobial specimen. From diagnostic perspective, such concentrations may not alter antibiotic susceptibilities of the isolated organisms as recently observed in urea (Iwalokun et al., 2003).

The components lipopolysaccharide and capsular polysaccharide in cell matrix constitute the carbohydrate requirements for almost all swarming pathogens including Proteus spp. (Consterton, 1987; Consterton et al., 1999). Mutant Myxococcus xanthus strains lacking O-LPS gene expression also lacked the ability to swarm (Bowden and Kaplan, 1998), while glycogen synthesis and swarming motility deficiencies have been observed in Escherichia coli deficient in CsrA, a pleitropic RNA - binding protein gene (Romeo et al., 1993; Wei et al., 2001). Gygi et al. (1995) also reported the involvement of a surface polysaccharide in swarming motility in Proteus mirabilis, similar to the spontaneous release of lipopolysaccharide by pathogenic P. aeruginosa (Cadieux et al., 1983). Our observed reduction in carbohydrate levels among the 40 treated Proteus strains seems to validate these findings.

Swarming motility, as a quorum sensing behavior, has been demonstrated in several signal transduction experiments to involve many distinct genes that promote intercellular communication (Fugua et al., 2001). synthesis of biosurfactants (Eberl et al., 1999) and virulence factors (Giskov et al., 1997). These findings have subsequently been validated by the works of Belas et al. (1998) and Duffour et al. (1998) where transposons mutants were employed to establish swarming motility virulence mediated gene expressions. and The localization of swarming and virulence genes within transposons implies that these genetic determinants are mobilizable (Duffour et al., 1998). Plasmids are extrachromosomal genetic elements that have also been implicated in the spread of resistance and virulence across species of pathogens through mobilization (Gebre-Yohannes and Drasar, 1988; Hale et al., 1984). The persistence of swarming motility behaviour in some of the Proteus strains after curing strongly suggests likely

the involvement of chromosomal DNA in swarming. While the observed unique plasmids of size 6.0, 12.5, 20 and 33.5 kb further implies the involvement of plasmids in this motility behaviour in *Proteus*. Therefore, studies aimed at defining the roles, regulation of plasmids and chromosomes associated with swarming in *Proteus* and other related organisms may further provide an insight on the dynamics of swarming motility behaviour and gene expression.

Based on the data obtained from this study, it can be concluded that swarming motility in *Proteus* may be coupled to the expression of plasmid genes, and is inhibited by SDS, urea and Tris *in vitro*.

ACKNOWLEDGEMENT

We are grateful to the laboratory technologists of Genetic Division of the Nigerian Institute of Medical Research (NIMR) for their assistance.

REFERENCES

- Adeyemo, AA; Gbadegesin, RA; Onyemenem, TN; Ekwezor, CC (1994). Urinary tract pathogens and antimicrobial sensitivity patterns in children in Ibadan, Nigeria. Ann Trop. Paediatr.14: 271 274.
- Alwen, J; Smith, DG (1967). A medium to suppress the swarming of *Proteus species*. J. Appl. Bacteriol. 30: 389 394.
- Belas, R; Schneider, M and Melch, M (1998). Characterization of *Proteus mirabilis* precocious swarming mutants: identification of rsbA, encoding a regulator of swarming behavior. J Bacteriol. 180: 6126 – 6139.
- Bowden, MG and Kaplan, HB (1998). The *Myxococcus xanthus* lipopolysaccharide O-antigen is required for social motility and multicellular development. Mol Microbiol. 30: 275 284.
- Cadieux, JE; Kuzio, J; Milazzo, FH anf Kropinski, AM (1983). Spontaneous release of lipopolysaccharide by *Pseudomonas aeruginosa.* J Bacteriol. 155: 817 – 825.
- Campanac, C; Pineau, L; Payard, A; Baziard-Mopuysset, G; Roques, C (2002). Interaction between biocide cationic agents and bacterial biofilms. Antimicrob. Agents. Chemother.46: 1469 – 1474.
- Costerton, JW; Cheng, KJ; Geesey, GG; Ladd, TL; Nickel, JC; Dasgupta, M and Marrie, TJ (1987). Bacterial biofilms in nature and disease. Annu. Rev. Microbiol. 49: 711 745.
- Costerton, JW; Stewart, PS and Greenberg, EP (1999). Bacterial biofilms: a common cause of persistent infections. Science. 284: 1318 1322.
- Cowan, ST (1974). Manual for identification of medical bacteria. 2nd ed. London. Cambridge University Press.
- Cutner, A; Cardoso, LD; Benness, CJ (1991). Assessment of urinary symptoms in early pregnancy. Br. J. Obstet Gynaecology. 98: 1283 1286
- Davies, DG; Parsek, MR; Pearson, JP; Iglewski, BH; Costerton, JW; Greenberg, EP (1998). The involvement of cell -to-cell signals in the development of a bacterial biofilm. Science. 280 : 295 298.
- Duffour, A; Furness, RB and Hughes, C (1998). Novel genes that upregulate the *Proteus mirabilis* flhDC master operon controlling flagellar biogenesis and swarming. Mol Microbiol. 29: 741 751.
- Eberl, Lp; Molin, S and Givskov, M (1999). Surface motility of Serratia liquifacuiens MG1. J Bacteriol. 181: 1703 1712.
- Fuqua, C; Parsek, MR; Greenberg, EP (2001). Regulation ofgene expression by cell-to-cell communication:Acyl-homoserine Lactone quorum sensing. Annu. Rev. Genet. 35: 439 – 468.
- Garriga, M; Hugas, M; Aymerich, T and Monfort, JM (1993). Bacteriocinogenic activity of lactobacilli from fermented sausages. J. Appl. Bacteriol. 75: 142 –148.

Gebre-Yohannes, A and Drasar, BS (1988). Transferable or mobilisable antibiotic resistance in *Shigella dysenteriae* types 1, 2, 3, 6 and 7 isolated in Ethiopia during 1974 – 85. J Med Microbiol. 27: 285 – 289.

Givskov, M; Eberl, L and Molin, S (1997). Control of exoenzyme production, motility and cell differentiation in *Serratia liquifaciens*. FEMS Microbiol Lett. 148: 115 – 122.

Greene, RV; Cotta, MA; Griffin, HL (1989). A novel symbiotic bacterium isolated from marine shipworm secretes proteolytic activity. Curr. Microbiol.19: 353 – 356.

Gygi, D; Rahman, MM; Lai, HC; Carlson, R; Guard-Petter, J and Hughes, C (1995). A cell surface polysaccharide that facilitated swarm cells of *Proteus mirabilis*. Mol Microbiol. 17: 1167 – 1175.

Hale, TL; Guerry, P and Seid, SC (1984). Expression of lipolysaccharide O-antigen in *Escherichia coli* K12 hybrids containing plasmids and chromosomal genes from *Shigella dysenteriae* I. Infect Immun.46: 472 – 475

Harshey, RM (1994). Bees aren't the only ones: swarming in Gramnegative bacteria.

Mol Microbiol. 13: 389 – 39

Hernandez, E; Cavallo, JD (1999). Abolition of swarming of Proteus. J Clin Microbiol. 37: 3435 – 3436

Iwalokun, BA and Akinwunmi, AO (2002). Swarming modulatory effects of some amino acids on *Proteus* strains from Lagos, Nigeria. Afr J Biotech. 1: 5 – 11.

Iwalokun, BA; Akinsinde, KA; Nkiruika, N (2003). Inhibition of swarming by urea and its diagnostic implications among uropathogenic *Proteus* species from Lagos, Nigeria. Afr J Clin Exp Microbiol. 4: 17 – 27

Jayaraman, J (1988). Laboratory manual in Biochemistry. 3rd edition. New Delhi. Wiley Eastern Limited. pp. 114 – 115, 130 – 131.

Kesah, CN; coker, AO; alabi, SA; Olukoya, DK (1996). Prevalence, antimicrobial properties and beta-lactamase production of haemolytic enterobacteria in patients with urinary tract infections in Lagos, Nigeria. Centr Afr J Med. 42: 147 – 150.

Liaw, SJ; Lai, HC; ho, SW; Luh, KT and Wang, WB (2000). Inhibition of virulence factor expression and swarming differentiation in *Proteus mirabilis* by p-nitrophenylglycerol. J Med Microbiol. 49: 725 – 731.

Liaw, SJ; Lai, HC; ho, SW; Luh, KT and Wang, WB (2001). Characterisation of p-nitrophenylglycerol- resistant *Proteus mirabilis* super-swarming mutants. J Med Microbiol. 50: 1039 – 1048

Liaw, SJ; Lai, HC; Ho, SW; Luh, KT and Wang, WB (2003). Role of RsmA in the regulation of swarming motility and virulence factor expression in *Propteus mirabilis*. J Med Microbiol. 52: 19 - 28

Lowry, OH; Rosebrough, NJ; Farr, AL; Randall, RJ (1951).Protein measurement with the folin phenol reagent. J. Biol Chem. 193:265 – 275. Mireles JR, Toguchi A, Harshey RM (2001). Salmonella enterica Serovar Typhimurium swarming mutants with altered biofilm –forming abilities: surfactin inhibits biofilm biofilm formation. J. Bacteriol. 183: 5848 – 5854.

Orret, FA (1999). Prevalence of Proteus species in urinary tract infections in a regional hospital in Trinidad. Zhonghua Yi Xue Za Zhi (Tapei). 62: 438 – 442.

Rasmussen, TB; Manefield, M; Andersen, JB; Eberl, L; Anthoni, U; Christophersen, C; Steinberg, P; Kjelleberg, S and Gbivskov, M (2000). How *Delisea pulchra* furanones affect quorum sensing and swarming motility in *Serratia liquifaciens* MG1. Microbiology.146: 3237 – 3244.

Reynolds DJ; Penn, CW (1994). Characteristics of *Helicobacter pylori* growth in defined medium and determination of its amino acid requirements. Microbiology. 140: 2649 – 2656.

Romeo, T; Gong, M; Liu, MY and Brun-Zinkernagel, AM (1993). Identification and molecular characterization of crsA, a pleiotropic gene from *Escherichia coli* that affects glycogen biosynthesis, gluconeogenesis, cell size and surface properties. J Bacteriol.175: 4744 – 4755.

Takahashi, S and Nagano Y (1984) . Rapid procedure for isolation of plasmid DNA and application to epidermiologic analysis. J. Clin Microbiol. 20: 608-615.

VanAsten, FJA; Gaastra, W (1999). Urea refrains swarming of *Proteus* mirabilis. J Clin Mirobiol. 37: 1652 - 1652

Wharton, DC and McCarty, RE (1972). Experiments and methods in Biochemistry. Macmillan, New York.

Wei, BL; Brun-Zinkernagel, AM; Simecka, JW; Prub, BM; Babitzke, P and Romeo, T (2001). Positive regulation of motility and flhDC expression by the RNA-binding protein CsrA of *Escherichia coli*. Mol Microbiol. 40: 245 – 256.

Williams, FD (1973). Abolition of swarming of *Proteus* by pnitrophenylglycerin: application to blood agar. Appl Microbiol. 25: 751 - 754.

Williams, FD and Schwarzhoff, RH (1978). Nature of swarming phenomenon in *Proteus*. Ann. Rev. Microbiol. 32: 101 - 122

Young, GM; Smith, MJ; Minnich, SA and Miller, VL (1999). The Yersinia enterocolitica motility master regulatory operon, flhDC, is required for flagellin production, swimming motility and swarming motility. J Bacteriol. 181: 2823 – 2833.