

African Journal of Fisheries Science ISSN 2375-0715 Vol. 6 (1), pp. 001-006, January, 2018. Available online at www.internationalscholarsjournals.org © International Scholars Journals

Author(s) retain the copyright of this article.

Full Length Research Paper

# Digestibility coefficients of processed jackbean meal *Cannavalia ensiformis* (L.) DC for Nile tilapia, *Oreochromis niloticus* (Linnaeus, 1758) diets

Jimoh W. A.<sup>1</sup>\*, Fagbenro O. A.<sup>2</sup> and Adeparusi E. O.<sup>2</sup>

<sup>1</sup>Fisheries and Aquaculture Unit, Department of Biological Sciences, Crescent University, Abeokuta, Ogun State, Nigeria.

<sup>2</sup>Department of Fisheries and Aquaculture Technology, Federal University of Technology, Akure, Ondo State, Nigeria.

# Accepted 5 October, 2017

A study was conducted to determine *in-vivo* apparent digestibility coefficient of nutrients in processed jackbean meal. Acid Insoluble Ash (AIA) was used as an indicator. The test diets consist of the seed meal replacing soybean meal at 10, 20, 30 and 40% to serve as test diets D10, D20, D30, and D40 respectively. A diet called as diet CTR, without jackbean meal served as control. These diets were fed to Tilapia *Oreochromis niloticus* fingerlings ( $6.37 \pm 0.07$  g) in a glass tanks for 5 weeks. Faecal collection was made by siphoning after four hours of each feeding. Proximate analysis of both diets and faecal samples were carried out for crude protein, lipid, crude fiber, energy, and AIA. These were used to calculate the digestibility coefficient of nutrients. There was no significant difference (p > 0.05) in lipid and organic matter digestibility of the fish fed control diet and test diets. Significant (p < 0.05) variations occurred in nitrogen-free extracts, protein and crude fiber digestibility of fish fed control diet and the test diets.

Key words: Digestibility, in vivo, jackbean, Oreochromis niloticus.

# INTRODUCTION

Soybean meal has high protein content and the best protein quality among plant protein feedstuffs used in fish feeds (Davies et al., 1999). It has been reported to partially or totally replace fishmeal in diets of many aquaculture species (Lovell, 1988; Lim and Akinyama, 1992). However, due to various other uses, the seed meal is now scarce and even expensive, beyond the reach of fish farmer. It then becomes a priority to look for alternative source. cheaper. protein Jackbean, *Cannavalia ensiformis* (L.) DC, an indigenous legume has shown potential as protein and energy sources of fish diets (Oliveira-Novoa, 1988) because of its high seed yield, energy and protein content. It has up to 30% crude protein and 60% carbohydrate (Udedibie, 1990). Digestibility of nutrients in fish diets needs to be studied because it is the digested feed, which is absorbed, that is

\*Corresponding author. E-mail: jawabus@gmail.com.

made available for cellular metabolism. The resultant of which will be tissue synthesis, and repair of worn-out tissues and various energy utilization channels (Yudkin, 1985; NRC, 1993). The most important characteristics of feedstuffs are the bioavailability of nutrients; hence reliable data on different ingredients for each species need to be well considered as a necessary prerequisite (Jauncey, 1993; Fagbenro et al., 2003). Borghesi et al. (2008) reported that knowing nutrient digestibilities of feed ingredients elicit interchangeability of feed ingredients without reducing animal performance. De Silva and Anderson, (1995) also observed that it is essential to have a knowledge of the digestibility of the main ingredients as well as of the whole diet in feed formulation and manufacture; In combination, chemical analysis and apparent digestibility coefficient (ADC) results allow us to precisely estimate not only the contribution of a particular protein source to a complete fish feed but also how much feed wastes and undigested nutrients (faeces) will potentially accumulate in fish pond (Koprucu and Ozdemer

Feed ingredient	Control	Test diets					
	CTR	D10	D20	D30	D40		
Fish meal	24.00	24.00	24.00	24.00	24.00		
Jackbean meal		6.00	12.00	18.00	24.00		
Soyabean meal	32.00	30.00	26.00	22.00	19.00		
Maize	24.00	22.00	20.00	18.00	15.00		
Cod-liver oil	2.50	2.50	2.50	2.50	2.50		
Vegetable oil	3.50	3.50	3.50	3.50	3.50		
Vit/min premix	2.00	2.00	2.00	2.00	2.00		
Cassava Starch	10.00	10.00	10.00	10.00	10.00		

Table 1. Gross composition (g/100 g dry matter) of experimental diets fed to *O. niloticus* at varying replacement levels of Jackbean meal.

2005). Various direct and indirect methods have been used to study digestibility. The direct method involves measuring all the feed consumed by the fish and all of the resulting excreta. A measured amount of feed and the faeces collected are analyzed for their nutrient content. The amounts of the nutrients in the excrements are then subtracted directly from those in the feed to determine the amounts retained (NRC, 1993). Smith (1971, 1976) and Smith et al. (1980) used an aquatic modification of the metabolism chamber designed for terrestrial animal studies; it allowed for the separate quantitative collection of gill, urine, and faecal excretions of rainbow trout. The fish were force-fed a measured amount of feed, and the various excrements were subsequently collected and analyzed for their nutrient content. The indirect method involves the use of a non-digestible exogenous or endogenous marker/indicator. It is assumed that the amount of marker in the feed and faeces remains constant throughout the experimental period and that all of the ingested markers appear in the faeces. The digestibility of the nutrient in question can then be determined by assessing the difference between the feed and faecal concentrations of the marker and the nutrient (NRC, 1993). Digestion coefficients determined by the indirect method have been useful and feeding regimens based on these data have been successful (Takeuchi et al., 1979; Cho and Kaushik, 1985; Wilson and Poe, 1985; Mangalik, 1986; Satoh et al., 1992). Fish can be sacrificed and the faecal samples either (i) removed from the lower large intestine (Smith and Lovell, 1971), or by (ii) gentle pressure in the abdomen of the fish (Nose, 1960a, b), or by (iii) suction method can be used (Windell et al., 1978). Faecal samples can also be periodically removed from the tank by siphoning or using settling

removed from the tank by siphoning or using settling column (Smith, 1989). Cho and Slinger (1979) have shown that if excretions are removed from the fish tank soon after expulsion, the collection of passively excreted faeces can give good digestibility data. Hence an attempt was made to evaluate *in vivo* apparent digestibility coefficient of nutrients in processed jackbean meal using acid insoluble ash as indicator.

### MATERIALS AND METHODS

#### Source of C. ensiformis seeds

The jack bean seeds were obtained from the National Root Crops Research Institute (NRCRI) in Vom, Plateau State, Nigeria. They were broken into coarse pieces (about 3 to 6 parts) using a grinding machine. The broken seeds were subjected to one hour ordinary cooking and the products was dried in the sun (30 to 32°C) for three days, milled and analysed for its proximate composition.

Five isonitrogenous diets (30% crude protein) were formulated as depicted in Table 1. The soybean meal protein in the diets was substituted at a rate of 10, 20, 30, 40% to serve as test diets designated as D10, D20, D30, and, D40 respectively. A diet, called as diet CTR, with soybean meal and without jack bean meal served as a control. The feedstuff was milled using grinding machine. Hot water was added to aid binding and then introduced into Hobart A-200T pelleting and mixing machine to obtain a homogenous mixture and then transferred through a mincer to produce 0.6 mm (long) 0.2 mm (diameters) pellets which were immediate sun dried (30 to 32°C). After drying for 3 days, the diets were kept in a freezer (-20°C).

#### **Culture condition**

Nile tilapia (Oreochromis niloticus) fingerlings ( $6.36 \pm 0.07$  g) were stocked in 15 glass aguarium tanks at the Department of Fisheries and Wildlife, Federal University of Technology, Akure, Nigeria. Fish were acclimated to laboratory conditions for seven days, during which they were fed a 30% crude protein diet before the experiment started. Fish were not fed for 24 h before the onset of the experiment. Triplicate batches of fish were fed to satiation twice a day (08:00 to 09:00 and 16:00 to 17:00 h) on each of the five experimental diets for an initial seven days before fecal collection began. Collection of faeces samples were carried out for 14 days by siphoning using a pipe (2 cm diameter) three hours after feeding. Uneaten diet was siphoned out 20 min after feeding. Droppings from the same tank were pooled together in a bowl and stored in freezer. The amount of dissolved oxygen (using a YSI model 57 oxygen meter) water temperature and pH (using electronic pH meter) were measured daily in all tanks.

#### **Proximate analysis**

The proximate analysis of the protein feedstuffs, diets and faecal matter samples were carried out in three replicates using the procedures described by the AOAC (1990). Moisture content was

Feed ingredient	Control	Test diets				
	CTR	D10	D20	D30	D40	
Protein	30.18	30.19	30.24	30.55	30.88	
Lipid	10.39	10.07	9.76	10.52	10.41	
Fiber	5.58	5.62	5.65	5.7	5.69	
Ash	6.04	5.09	5.41	5.26	5.3	
NFE	39.56	40.08	40.84	39.61	38.96	
Moisture	8.25	8.95	8.1	8.36	8.76	
Energy (Kcal /Kg)	430.89	430.06	430.53	431.42	432.83	
AIA (%)	3.05	3.18	3.08	3.11	3.36	

 Table 2. Proximate composition (%) of experimental diets fed to O. niloticus at varying replacement levels of Jackbean meal.

NFE-Nitrogen free extract, AIA- Acid insoluble ash.

determined as a percentage loss in weight after drying a known weight of the sample at 100°C until constant weight was obtained. Crude protein was determined by digesting a known weight of the sample ( $w_1$ ) in Kjedahl flask unit using 25 ml of concentrated H<sub>2</sub>SO<sub>4</sub> and a gram of copper catalyst. The percentage nitrogen was then calculated as;

% Total Nitrogen (N) = 
$$\frac{T_1 \times M_2 \times 0.014 \times V}{W_1 \times V_2} \times 100$$

T = Titre Value Ma = Molarity of acid W1 = Weight of the sample V1 = Initial Volume put in a distillation unit V2 = Final volume Obtained from the distillation % Crude Protein = N x 5.4 (Legume)

A factor value of 6.25 was used to convert nitrogen to protein for animal protein. The lipid content was determined by subjecting the sample to a continuous extraction with petroleum ether using Soxhilet apparatus. The residue from ether extraction was subjected to successive treatment of boiling with dilute acid and dilute base, after which it was filtered and ignited in a furnace. The difference obtained between the burnt and unburnt fractions was calculated to be crude fibre. Ash Content was determined by subjecting the already dried samples with known weight to ignition in a muffle furnace at 550°C for 8 h. The nitrogen free extract was estimated by difference. Energy contents of faeces and the diets were determined using an adiabatic bomb calorimeter.

#### Acid insoluble ash (AIA) analysis

AIA analyses were carried out on the diets and feces. AIA was obtained by adding 25 ml of 10% HCl to the weighed ash content of a sample. This was covered with a water-glass and boiled gently over a low flame for five minutes. This was then filtered using ashless filters and washed with hot distilled water. The residue from the filter was returned to the crucible and ignited until it was carbon free after which it was weighed. Percentage AIA was calculated as;

$$%AIA = \frac{Weight of AIA}{Weight of Ash} \times 100$$

#### Determination of digestibility coefficient

This was calculated on the percentage of AIA in feed and in faeces and the percentage of nutrient on diets and feaces

Apparent Organic matter Digestik	oility (%)	AIA in Diets
	- 100	AIA in Faeces
Apparent Digestibility (%) $_{=100}$	(AIA in Diets)	×(Nutrients in Faeces)
	(AIA in Faeces)	(Nutrients in Diets)

#### Statistical analysis

All determinations were conducted in triplicates and the means were subjected to Analysis of Variance, where, the ANOVA revealed a significant difference, Fishers Least Significant Difference (LSD) was used to compare differences among individual treatment means using SPSS version 13.

## **RESULTS AND DISCUSSION**

The ingredient and proximate composition of experimental diets is presented in Tables 1 and 2 respectively. The result revealed the diets were isonitrogenous and isocaloric conforming to the recommendation of optimum dietary protein of the fish (Luquet, 1991). The lipid requirement of the fish also met the standard of the aquadiets (Jauncey and Ross, 1982). Acid Insoluble Ash (AIA) was used as a marker because it is more reliable indicator of digestibility coefficient (Halver et al., 1993) since the dietary ingredient (ash) is used and analysis of this component in feces collected uses simple gravimetric technique. Defecation of the fish started about four hours after feeding and lasted for 15 to 25 min. Same observation was made by Adeparusi and Jimoh (2002).

Table 3 revealed the proximate composition of the faecal sample. A general reduction in the nutrients of the faeces as compared to the diets showed that some percentages of nutrients were absorbed and made available for metabolism. There was a significant

Demonstern	Control	Test diets				
Parameter	CTR	D10	D20	D30	D40	
Protein	8.82 <sup>b</sup>	10.64 <sup>b</sup>	10.97 <sup>b</sup>	12.01 <sup>ab</sup>	14.01 <sup>a</sup>	
Lipid	3.11	3.02	3.05	2.74	3.3	
Fibre	3.92 <sup>a</sup>	4.1 <sup>a</sup>	4.16 <sup>a</sup>	4.31 <sup>a</sup>	4.48 <sup>b</sup>	
Ash	12.4 <sup>a</sup>	14.04 <sup>a</sup>	14.11 <sup>a</sup>	14.12 <sup>a</sup>	13.98 <sup>b</sup>	
NFE	60.8	60.19	57.8	56.94	54.18	
Moisture	10.95	8.01	9.91	9.88	10.05	
Energy Kcal /Kg	328.5	333.54	327.78	327.2	332.48	
AIA (%)	8.9	8.95	9.78	9.12	8.94	

 Table 3. Proximate composition (%) of faecal samples of O. niloticus fed varying replacement levels of Jackbean meal.

Table 4. Apparent digestibility coefficients of the nutrients in the diets fed to O. niloticus.

<b>D</b>	Control	Test Diets				
Parameter	CTR	D10	D20	D30	D40	
Crude protein	89.98 <sup>a</sup>	87.48 <sup>ab</sup>	88.57 <sup>ab</sup>	86.59 <sup>b</sup>	82.96 <sup>c</sup>	
Crude fibre	74.88 <sup>a</sup>	74.69 <sup>a</sup>	76.79 <sup>a</sup>	74.26 <sup>a</sup>	70.38 <sup>b</sup>	
Lipid	89.7	89.28	90.08	91.13	87.97	
NFE	55.36 <sup>a</sup>	50.98 <sup>0</sup>	46.62 <sup>c</sup>	46.65 <sup>c</sup>	47.77 <sup>c</sup>	
Energy	76.02 <sup>a</sup>	74.32 <sup>b</sup>	73.87 <sup>b</sup>	72.28 <sup>c</sup>	71.12 <sup>c</sup>	
Organic matter	64.67	63.26	67.33	65.52	61.53	

difference (p < 0.05) in the crude protein, ash and nitrogen free extract of the faecal samples tested. Diet D4O had the highest fecal nitrogen. The result agreed with Adeparusi and Jimoh (2002) that there was a reduction in nutrient value of faecal sample of *O. niloticus* fed lima bean (*Phaseolus lunatus*) when compared to the diets.

Table 4 presented the apparent digestibility coefficient of nutrients in the diets fed to *O. niloticus* fingerlings. A significant difference (p < 0.05) was observed between different nutrients in the different diets. The results of the apparent protein digestibility of the seedmeal are similar to that reported in Fagbenro (1988) for the same seed meal fed to *O. niloticus* but lower than the values reported by Martinez-Palacios et al. (1988) who fed the same seed meal to *O. mossambicus*. The little variation observed could be attributed to variability of nutrients as well as differences in nutrient processing, experimental methodology and feces sampling technique (Jauncey, 1993).

No significant difference (p > 0.05) occurred for crude fibre digestibility of *O. niloticus* fed jackbean meal at varying replacement level except for diet D40; the crude fibre digestibility coefficient in this study was higher than the carbohydrate digestibility in the same experiment. This could be attributed to the natural feeding habit of tilapia that consists mainly of plant material (Pullin, 1983).

O. aureus was found to digest highly fibrous feedstuffs such as alfalfa meal (Mgbenka and Lovell, 1987). Crack and cook processing method improved the digestibility of crude fibre. This result conforms to the report of Adeparusi and Jimoh (2002) that fibre digestibility of O, niloticus fed lima bean diet was improved with toasting and autoclaving. No significant (p < 0.05) variation was observed in the lipid digestibility of O. niloticus fed with processed jackbean meal. The high lipid digestibility by this species was found to be in line with what was reported in Hossain et al. (1992) for rainbow trout. A range of 76 to 97% fat digestibility of various sources of fat has been reported for channel catfish (Lovell, 1977). Andrew et al. (1978) reported that the ability to digest fat appears to be influenced by temperature and the level of fat in the diet.

There was a significant difference (p < 0.05) in the carbohydrate digestibility of *O. niloticus* fed processed jackbean meals and the control, however there was no significant difference (p > 0.05) in the carbohydrate digestibility of the fish fed diet CTR, DIO and D40. The low carbohydrate digestibility recorded in this study was similar to that reported by Adeparusi and Jimoh (2002) for *O. niloticus* fed lima bean. The result however deviated from the report of Popma (1982) that warm water fish like *O. niloticus* absorbs 60% or more of raw corn starch. The digestibility of carbohydrates has been

Table 5. Mean value of water quality parameters in the experimental tanks.

Parameter	Control	Test diets			
	CTR	D10	D20	D30	D40
Temperature (°C)	26.20±0.21	26.40±0.73	26.58±0.32	26.10±0.15	26.35±0.32
Dissolved oxygen (g/dm <sup>3</sup> )	6.78±0.29	6.55±0.32	6.56±0.18	6.59±0.54	6.79±0.64
рН	7.11±0.35	6.98±0.21	7.20±0.11	6.92±0.53	7.04±0.27

shown to vary with their complexity, source treatment and level of inclusion in the diet (Phillip, 1972; Lovell, 1977; Cho and Slinger, 1979). Generally tilapia have been reported to have ability to digest carbohydrates relatively well in feedstuff but not as well as fat or protein Lovell (1988). There was significant difference (p < 0.05) in the energy digestibility of *O. niloticus* fed processed jackbean meals and the control. However, there was no significant difference (p > 0.05) in the energy digestibility of the fish fed with D20 and CTR and the fish fed with D10 and D40. The high apparent digestibility coefficient of energy recorded in this study is similar to the values recorded by Fagbenro (1998) of the same seed meal fed to *O. niloticus*. Lovell and Durve (1982) revealed that cooking improved gross energy digestibility.

These were no significant difference (p > 0.05) in the organic matter digestibility of *O. niloticus* fed processed jackbean meals and the control. The organic matter digestibility coefficient reported in this study is similar to that of Fagbenro (1998) and Olivera et al. (1988), but lower than the values reported by Martinez-Palacios et al. (1988). The variation may be attributed to processing methods and or experimental methodology. The crack and cook processing technique applied to jackbean meal improved its digestibility.

Table 5 revealed the physico-chemical conditions under which the experiment was conducted. The values were within the optimum range for the normal physiological functioning of not only tilapia but also most of the warm water fish species (Boyd, 1990). Lazo and Davies (2000) reported that experimental culture system must be suitably designed to allow maximum growth of fish and water provide adequate quality throughout the experiment. Conclusively, this study reveals that up to 20% of soybean meal in the diets of O. niloticus can be replaced by jackbean meal without negatively affecting the fish performance vis-à-vis the rate at which the nutrients are absorbed in the species. This will reduce the cost of fish feed considerably.

#### REFERENCES

- Adeparusi O,Jimoh WA (2002). Digestibility coefficients of raw and processed lima bean diet for Nile Tilapia, *Oreochromis niloticus*. J. of Appl. Aquac., 1-2(3): 89-98.
- Andrew JW Murray, MW Davies JM (1978). The influence of dietary fat levels and environmental temperature on digestible energy and absorbability of animal fat in channel catfish. J. Nutr. 108: 749-752.

AOAC (1990). Association of official Analytical chemistry. Official method of analysis, 11th edition: Washington DC.

- Borghesi R, Portz L, Oetterer M, Cyrino JEP (2008). Apparent digestibility coefficient of protein and amino acids of acid, biological and enzymatic silage for Nile Tilapia (*Oreochromis niloticus*). Aquaculture Nutr., 14: 242 248.
- Boyd CE (1990). Water Quality in warm water fish pond, Auburn University Agriculture Experiment Station, Auburn Al.
- Cho EY, Ślinger SL (1979). Apparent digestibility measurement in foodstuff for rainbow trout, in J.O. Halver and K. Tiews Eds. World Symposium on Fin Fish Nutrition and Fish Feed Technology Heinemann, Berlin. Germany. 2: 239 247
- Cho CY, SJ Kaushik (1985). Effects of protein intake on metabolizable and net energy values of fish diets. Pp. 95-11 in Nutrition and Feeding of Fish, C. B. Cowey, A. M. Mackie, and J. G. Bell, eds. New York: Academic Press.
- Davies S, Fagbenro OA, Abdelwaritho A, Diller I (1999). Use of soybean products as fishmeal substitute in African catfish, *Clarias gariepinus*. Appl. Trop. Agric., 14(1): 10-19.
- De Silva SS, Anderson TA (1995). Fish Nutrition in Aquaculture Chapman and Hall, London.
- Fagbenro AO (1998), Apparent digestibility of various legume seed meals in Nile tilapia diets short communication. Aquac. Int., 6: 83-87.
- Fagbenro OA, Akande TT, Fapounda OO (2003). Use of Roselle (*Hibiscus Sabdariffa*) seed meal as a soybean meal replacer in practical diets for fingerlings of African Catfish (*Clarias gariepinus*). Proceeding of the third international conference on African fish and Fisheries, Cotonou, Benin, 10-14 November, Ed J. Snoeks, P. Laleye and P. Vandewalle pp. 73-79.
- Halver JC, Yiman A, Smith RR (1993). Acid Insoluble Ash as a convenient method for estimating digestible components in diet. Pg 230 in proceeding international conference world on Aquaculture and fresh water fish.
- Hossain MA, Nahar N, Kamal M, Islam MN (1992). Nutrient digestibility coefficient of some plants and animal proteins for tilapia (*Oreochromis niloticus*) J. Aquaculture Trop., 7: 257-260.
- Jauncey K (1993). Advances in freshwater fish nutrition. In Feed Production Tomorrow. Victam International Leiden, The Netherlands pp. 16-41.
- Jauncey K, Ross B (1982). A Guide to Tilapia feeds and feeding.
- Institute of Aquaculture, University of stirling, Scotland. pp 3. Köprücü K, Özdemir Y (2005). Apparent digestibility of selected feed ingredients for Nile Tilapia (*Oreochromis niloticus*). Aquaculture, 250:
- 308 316. Lazo JP, Davies DA (2000). Ingredients and feed evaluation in Encyclopedia of Aquaculture edited by R.R. Stickney. A Wiley Interscience Publication, John Wiley and Sons incorporation, USA, pp 1063.
- Lim C, Akinyama DM (1992). Full fat soybean utilization by fish. Asian Fisheries Sci., 5: 181-197.
- Lovell RT (1977). Digestibility of nutrients in feedstuff of Catfish. In nutrition and feeding of channel catfish by R.R. Stickney, R.T. Lovell (eds) southern cooperative ser. Bull. 218. Auburn, Ala, Auburn University.
- Lovell RT (1988). Use of soybean product in diets for Aquaculture species. J. Aquat. Prod., 2: 27-52.
- Lovell RT, Durve VS (1982). Effect of grinding and heating on digestion of gross energy in feedstuff for channel Catfish. Fisheries Ann. Rep. 1979, Auburn University, Auburn Al.

- Luquet P (1991). Tilapia (*Oreochromis Spp*) in R.P Wilson (ed). CRC handbook of fin fish. CRC press, Inc. florida. p. 161-179
- Mangalik A (1986). Dietary energy requirements of channel catfish. Ph.D. dissertation. Auburn University, Auburn, Alabama.
- Martinez-palacios CA, Galvan Cruz R, Olvera Novoa MA, Chavezmartinez C (1988). The use of Jackbean (*canavalia ensiformis* Leguminosae) meal as a partial substitute for fish meal in diet for tilapia (*Oreochromis mossambicus* Cichlidae). Aquaculture, 68: 165-175.
- Mgbenka BA, Lovell RT (1987). Digestibility of feedstuff and supplementary diet by grass carp. Nigerian J. Appl. Fish. Hydrobiol., 2: 65-71.
- Nose T (1960a). On the digestion of food protein by goldfish (*Carassius auratus* L.) and rainbow trout (*Salmo gairdneri* G). Bull. Freshwater Fish. Res. Lab., 10(1): 12-22.
- Nose T (1960b). On the effective value of freshwater green algae, *Chlorella ellipsoidea*, as a nutritive source to goldfish. Bull. Freshwater Fish. Res. Lab., 10(1): 1-10.
- NRC (1983). Nutrient requirements of warmwater fishes and shellfishes. National Academic press Washington DC. pp. 102.
- NRC (1993).Nutrient requirements of fishes. National Academic press Washington DC. pp. 100.
- Olvera-Novoa MA, Martinez PCA, Galvan CR, Chavez, SC (1988). The use of seed of the leguminous plant sesbania grandiflora as a partial replacement for fishmeal in diets of tilapia (*Oreochromis* mossambicus). Aquaculture, 71: 51-60.
- Phillip AM (1972). Caloric and energy measurement in fish Nutrition (J.E. Halver eds), New York and London Academic press.
- Popma TJ (1982). Digestibility of selected feedstuffs and naturally occurring algae by Tilapia Ph.D. Dissertation, Auburn University, Auburn Al.
- Pullin RSV (1983). Choice of tilapia species for aquaculture. Page 87-114 in Pullin R.S.V and R.H.L McConnell eds. Proceeding International Conference on Tilapia in Aquaculture, Tel Aviv, Israel.

- Satoh S, CY Cho, T Watanabe (1992). Effect of fecal retrieval timing on digestibility of nutrients in rainbow trout diet with the Guelph and TUF faeces collection systems. Nippon Suisan Gakkaishi, 58: 1123-1127
- Smith RR (1989). Nutritional energetic. in fish nutrition (J.E Halver eds). New York Acaedemic Press. pp. 1-29
- Smith RR (1971). A method for determination of digestibility and metabolizable energy of feedstuffs for finish. Prog. Fish-Cult., 33: 132-134.
- Smith RR (1976). Metabolizable energy of feedstuffs for trout. Feedstuffs, 48: 16-21
- Smith BW, RT Lovell (1971). Digestibility of nutrients in semipurified rations by channel catfish in stainless troughs. Proc. Annu. Conf. Southeast Assoc. Game Fish Comm., 25: 425-459.
- Smith RR, MC Peterson, AC Allred (1980). The effect of leaching on apparent digestion coefficients in determining digestibility and metabolizable energy of feedstuffs for salmonids. Prog. Fish-Cult., 42: 195-199.
- Takeuchi T, T Watanabe, C Ogino (1979). Digestibility of hydrogenated fish oils in carp and rainbow trout. Bull. Jpn. Soc. Sci. Fish, 45: 1521-1525.
- Udedible ABI (1990). Nutitional evaluation of jackbean (*Canavalia ensiformis*) for the Nigerian Poultry Industry. AMBIO, 19: 361-365.
- Wilson RP, W E Poe (1985). Apparent digestibility of protein and energy in feed ingredients for channel catfish. Prog. Fish-Cult., 47: 154-158.
- Windell JT, JW Foltz, JA Sarokon (1978). Methods of faecal collection and nutrient leaching in digestibility studies. Prog. Fish-Cult., 40(2): 51-55
- Yudkin J (1985). The Penguin encyclopedia of nutrition, New York, Viking, pp. 121