

Full Length Research Paper

Does illiteracy influence pregnancy complications among women in the slums of greater Mumbai

Vijay M. Sarode

Mulund College of Commerce, Mulund (West), Mumbai-400 080, India. E-mail: vijaymsarode@yahoo.com.

Accepted 18 October, 2018

This paper examines utilization of health services available to the women in the slums on hilly area in Mumbai and also checks whether non utilization of antenatal care (ANC) and having reproductive health problems during pregnancy create complications during child delivery vis-à-vis standard of living index constructed from household amenities, housing quality, drinking water, electricity and toilet facilities. This study uses primary data collected using cluster sampling of a sample size of 346 reproductive women who have given at least one live birth prior to the survey on the education of the study women, antenatal care indicators, antenatal check-ups and reproductive health problems during pregnancy and complications during child delivery among the slum dwellers of Ramabai Nagar was studied. The findings using logistic regression reveal unimaginably low level of utilization of health services by illiterate women in the study area. Besides these, there is evidence that these respondents did not go for ANC and faced reproductive health problems during pregnancy which created problems during child delivery, particularly to illiterate mothers. This paper suggests that awareness is very much required at every stage of ANC particularly to illiterate women with low SLI category in the slums in order to ensure reproductive health during pregnancy.

Key words: ANC, pregnancy complications, illiteracy, Mumbai slums.

INTRODUCTION

Every seven minutes one woman dies due to complications in pregnancy or child birth in India (The Registrar-General of India, 2007). That makes 77,000 deaths every year. This is further compounded by the high percentage of deaths in the run-down maternity services and mother-and- childcare centers and rural health facilities. The current survey also reports that there are 300 maternal deaths per lakh (100,000) live births and what is most worrying is that women in the 20 - 24 age group constitute one-third of the total deaths. The main causes for a majority of deaths are haemorrhage, puerperal sepsis (infections after delivery), complications of abortion, obstructed labor, and hypertensive disorders associated with pregnancy and lack of education and awareness (Registrar General of India, 2006). Most of the women are anemic because they don't take good care of themselves. Inadequate health care and child birth in quick succession are the other issues. Abortions done by dais and quacks, which is widely prevalent in India, could also give rise to various infections.

In the poorest parts of the world, the risk of a woman dying as a result of pregnancy or childbirth during her

lifetime is about one in six compared to about one in 30,000 in Northern Europe. India being a developing country, contributes 26% of the global burden of maternal deaths with nearly 136,000 women dying annually (UNICEF, 2009) due to causes related to pregnancy and childbirth. Such a discrepancy poses a huge challenge in meeting the fifth Millennium Development Goal to reduce maternal mortality by 75% between 1990 and 2015 (Carine, 2006). It has also been estimated that for every maternal death, there are over 100 acute morbid episodes, indicating overall figure of 62 million maternal morbidities annually (Koblinsky, 1993). This invites the attention of researchers to examine the causes and epidemiological factors associated with maternal deaths.

Antenatal care provides a preventive service that monitors pregnancy for signs of complications, detects and treats pre-existing and concurrent problems of pregnancy, and provides advice and counseling on preventive care, diet during pregnancy, delivery care, postnatal care and related issues, thereby reducing maternal morbidity and mortality, if delivered effectively. However, the success that it has in achieving this aim is related to the quality

of service that is provided, the number of visits that a woman receives during pregnancy, the timing of those visits and the existence of and accessibility to professional delivery care when necessary (World Health Organization, 1996b).

In 1996, safe motherhood and child health services were incorporated into the Reproductive and Child Health (RCH) Programme in India. This program recommends that as a part of antenatal care, women receive two doses of tetanus toxoid vaccine, adequate amounts of iron and folic acid tablets (90+) or syrup to prevent and treat anaemia, and at least three antenatal check-ups that include blood pressure checks and other procedures to detect pregnancy complications (Harrison, 1990; Ministry of Health and Family Welfare, 1997, 1998b). Maharashtra (India) has the highest number of slum dwellers (Kapadia et al., 2002) having a slum population of more than 40 million (Census of India 2001). The study conducted (Godbole and Talwalkar, 1999) in urban slums in Maharashtra (excluding Greater Mumbai) showed that the antenatal care coverage for three or more ante-natal check-ups was 55 percent for women in the slums. About 58 per cent women in slums reported that they have taken a complete dose of iron and folic tablets and only 34 percent women reported a birth interval of more than three years in slum areas. Anemia is an underlying cause for a range of morbidities and severe anemia is a cause of maternal mortality. The consequences of anemia are severe, long term and often irreversible. A study (Khilare 2001) conducted in the slum area of Pimpri-Chinchwad area of Pune district indicated that out of a total of 1,797 women registered for antenatal care at the PCMC Bhosari hospital in 2000, about 83 percent were anemic ($hb < 11$ gms/dl). The proportion of anemic pregnant women increased to 89.6 percent for unrecognized slums.

The study conducted (Jean, 2008) in slum of Nairobi where access to appropriate facilities is limited, women with no formal education, availed less appropriate health facilities. Un-educated women, not seeking delivery care at a proper facility are more likely to have pregnancy complications in the later stage of pregnancy. Besides, women who delivered at a health facility indicated that more than 75% of women who delivered at appropriate facilities had at least one complication during delivery compared with about 66% among those who delivered at inappropriate facilities ($p < 0.01$).

In another study conducted in slum areas of Dhaka, Bangladesh, women who delivered in facilities (both elective and emergency transfers) were of significantly higher economic status, and better educated and significantly more likely to have received antenatal care than women who gave birth at home. There were no significant differences in parity, nutritional status, anemia, or serious delivery-related complications between those who delivered at home and those delivered at elected facility. There was no significant difference in postpartum morbidity caused by delivery location ($p = 0.58$).

Emergency transfers, however, were more likely than deliveries at home or at elective facility among "primi-para" (odds ratio [OR] = 1.9; $p < 0.01$) and among those who met the operational definitions for serious delivery-related complications (OR = 3.4; $p < 0.01$). This study also showed that the self-reported postpartum morbidity was associated with maternal characteristics, delivery-related complications and some birthing practices (Fronczak et al., 2007).

A study (Mayank et al., 2001) of women in the slums of Delhi, (India) reports that among the women who experienced bleeding during pregnancy, 44 percent actually recognized it as a danger signal; of those experiencing high blood pressure and swelling of the hands, face and feet, only 33 percent and 9 percent respectively, recognized its severity. Indeed, this study concludes that although the vast majority of pregnant women received antenatal care, fewer than 10 percent had been informed about danger signals.

In Andhra Pradesh, Madhya Pradesh and Orissa, the states of India, women who did not seek care for complications experienced during pregnancy cited lack of mobility and lack of resources as reasons while those who did not seek care for complications experienced in the postpartum stage cited lack of resources as the reason (Murthy and Barua, 2001).

Research study on Urban Female Sample Survey conducted by the Institute of Health Management, Pachod (IHMP) in 27 slums of Pune, (India) indicates that women suffer from much preventable morbidity. Post-abortion complications are reported in 42 percent of the cases. As many as 44 percent women from urban slums did not seek treatment for reproductive tract infections. Data also indicate that 68 percent women harbour negative gender attitudes against themselves, a result of the process of socialization. These attitudes have a direct impact on their treatment-seeking behaviour and utilization of antenatal services (Kapadia-Kundu and Tupe, 2001).

It has also been ascertained that pregnancy-related problems have far-reaching consequences on the overall reproductive health of women, in addition to their contribution to maternal mortality (Bhatia Cleland, 1995a). Besides less attention has been paid on pregnancy related problems and treatment to these problems. Gynecological morbidity has been studied at community level to a certain extent (Bang et al., 1989; Bhatia Cleland, 1995b; Parikh et al., 1996; Prakasam, 2004) but study on pregnancy related problems and complications during delivery are scanty. Recently, a few studies have been conducted mainly to explore the prevalence of obstetric morbidity at the community level (Bhatia et al., 1996; Srinivasa et al., 1997), but negligible attempts were made to examine the determinants of obstetric morbidity among women in slums. Women, particularly in the poor socio-economic status have the inherent tendency of late reporting of morbidity episodes, which subsequently



Figure 1. Profile of the study area.



Figure 2. Environmental condition in study area.

worsens the intensity of health disorders and co-morbidities (Harikrishnan, 2009).

In general, women in the slum remain unaware of their own reproductive health problems occurring during pregnancy such as danger sign for pregnancy, excessive bleeding, anaemia, diet care during pregnancy and blood pressure checks etc. Further risk involved in repeated pregnancies and proper utilization of antenatal and post-

natal care is another concern. Hence it is necessary to impart knowledge about pregnancy related problems and to understand the root cause of generating complications during delivery among such poor women in slums. Besides, women in the urban slums are unaware of the existing health facilities and even though these facilities are available, it has not been adequately utilized. Thus, keeping in view of the above research work, an attempt is made, to evolve a suitable strategy for knowing

- i) The reproductive health of the women during pregnancy,
- ii) utilization of antenatal care among study women studied and
- iii) the determinants influencing complications while delivery among women in slum in the area of Greater Mumbai.

Background of the study area

According to a UNESCO document (1997), "a slum is a building, a group of buildings, or area characterized by overcrowding, deterioration, insanitary conditions or absence of facilities, or amenities which, because of all these conditions or any one of them, endanger the health, safety or morals of its inhabitants or the community (Anderson, 1960). "Slums may be characterized as areas of substandard housing condition within a city (Nangia and Thorat, 2000) . A slum is always an area. A single, neglected building even in the worst stage of deterioration does not make a slum" (Bergel, 1955). Apart from these definitions, slum is an area of darkness, an area of poverty and thus poverty is the prime characteristic of slum.

Likewise, Ramabai Nagar, a densely populated slum, situated on the hilly area of Bhandup, comes under S ward of Brihan Mumbai Municipal Corporation, Mumbai and is about 4 km away from Bhandup railway station, a suburb in the Central Mumbai, India. This ward spreads over 25.22 sq km and has an overall population density of 27,398 per sq km (Census of India, 2001, Maharashtra population data with data on slum population in urban units). Bhandup is ranked second position as far as slum area populations in Mumbai are concerned. In Bhandup, out of 6.91 lacs of people, 5.71 lacs of people stay in the slum area. The slum sex ratio of S ward is 816 while female and male literacy rates are 81.67 and 93.02. This rate is above the national level.

Ramabai nagar slum was formed before 1980. This slum was constructed on the forest land. Most of the residents are either migrants or tribals. This slum area has two parts, Ramabai nagar no. 1 and 2 and spreads over 2 sq km. The present study was conducted in Ramabai nagar slum no. 2 (Figure 1-4) where the population of this slum was 3500. Many of the residents stay in semi-pucca houses having roof of asbestos sheets or tin or squared tiles. Homes either consist of huts




FIG 4 Lane in the study area.




Figure 3. Housing condition in the study area.

(zopadis) constructed largely of wooden planks, cane, bamboo and occasionally, brick and tiles or of single-room flats in concrete buildings called (chawls), each of which contains about 20 - 25 rooms. The area is very much congested and has no amenities worth mentioning. Drinking water facility in the study area is very poor. Many of them do not have their own water taps. Common

tap is provided in the area outside their home and they use it for washing clothes and utensils and even for bathing and this leads to the formation of open drains. Toilet facilities are inadequate with an average of one toilet for about 30 chawl residents. Narrow pathways crisscross the slums and are lined with open drains and open garbage. Smoke emitted by nearby factories makes the slum area highly air polluted. Besides, it has very limited open space. All these cause many chronic diseases and results in high rate of mortality. Most of the inhabitants are migrants from UP, Bihar, West Bengal and interior parts of Maharashtra. Such migrants have low income and so cannot afford to give good education and sufficient food to their wards. Accessing medical facility is another hurdle. The sub-center is situated at the bottom of this hilly area and walking down and reaching back home itself takes almost one hour. Community health volunteers provide essential medical facilities to the slum dwellers, particularly to the pregnant women during antenatal care (ANC) and postnatal care (PNC) periods.

For child delivery, the study women avail themselves of the government hospital facility near Bhandup railway station while many others prefer delivery at home. For ANC and PNC some of them do visit sub-center of urban health center. In short what is observed is that the life of these residents is very hard as they are deprived of pollution free environment, hygienic and sufficient food, pure drinking water, good education and health facility as they stay on hilly area.

MATERIALS AND METHODS Measuring

household standard of living

In the absence of data on income and consumption measures, household standard of living indices are often constructed using three sets of information, namely, source of drinking water, toilet facility and type of house and ownership of selected consumer durables (Montgomery et al., 2000). Index scores for the present study ranges from 1 - 6 for a low SLI to 7 - 9 for a medium SLI and ≥ 10 for a high SLI (Appendix). There are three other approaches in the construction of the standard of living indices differing in the manner in which different household amenities, quality of housing materials and assets are weighed.

Data

For the present investigation, two stage sampling procedures have been adopted. In the first stage, the slums in the Greater Mumbai according to the size of their population were listed using the "Directory of Slums" published by the Office of the Additional Collector (ENC), Mumbai and Mumbai Sub. Dist. Two lists were prepared, one for plain area slums and the other for hilly area slums. From each list, one slum was selected at random. The slum selected from hilly area was Ramabai Nagar slum in Bhandup (w). The population of this slum (study area) was approximately 3500.

In the second stage of sampling from the study areas using cluster sampling, two clusters were selected at random. From these two clusters, 349 households were selected. The respondents were interviewed carefully using structured schedule by trained




Figure 5. Utilization of antenatal care services showing problems during pregnancy in Ramabai Nagar Slum Hill, Bhandup, Mumbai, India.

investigators who generally work with the doctors in Pulse-Polio campaign. On the whole, this study covers 349 reproductive women in slums representing slum population in Greater Mumbai. This survey was conducted from June to August 2005. Before going for the survey, as pilot survey was also conducted.

In order to know the reproductive health of study women during pregnancy utilization of antenatal care, the determinants influencing complications during delivery and children born to the mothers during three years prior to the survey were considered.

Method of analysis

Logistic regression analysis was used to assess the effect of reproductive health problems during pregnancy with no ANC on complications during delivery controlling for other variables included in the model. For the logistic regression analysis purpose, the reproductive women who have given at least one live birth during three years prior to the survey were considered.

RESULTS AND DISCUSSION

Reproductive health problems during pregnancy

Figure 5 shows the problems during pregnancy reported by study women. About 47 percent of the women in the study area reported that they had at least one problem during pregnancy. The major antenatal problems reported were excessive fatigue (48 percent) (NFHS-2: 49.1 percent), followed by excessive vomiting (42 percent), swelling of the legs (31 percent) (NFHS-2: 35.9 percent), pain in abdomen (21 percent), white discharge (15 percent), blurred vision (10 percent) (NFHS-2: 12.1 percent), vaginal bleeding (9 percent) (NFHS-2: 3.5

percent), convulsion not from fever (7 percent) (NFHS-2: 10.5 percent), night blindness (6 percent) (NFHS-2 and RCH: 6.3 percent), and anemia (4 percent) (NFHS-2: 16.1 percent). The percentage of these pregnancy problems remains almost same as the problems specially shown in Mumbai slum data in NFHS-2 for Maharashtra state where the survey was taken in 1998 - 1999. This indicates that even after a period of 7 years, the reproductive health condition of study women living in slum on hill remains poor. Probably these women are not utilizing the medical facilities available in that area.

Antenatal care

Women not receiving antenatal check-ups tend to be disproportionately older women, women of high parity, women from scheduled tribes, illiterate women and women from households with low standard of living.

Figure 6 shows the reasons for not receiving an antenatal check -up according to residence in Ramabai nagar slum area, Bhandup. About 41.7 percent of mothers in the study area from Ramabai nagar reported that they did not go for a check-up due to fear. 16.7 percent of study women said that they had no money. 13.9 percent of study women said, they could not decide whether to go for antenatal check -up and 27.8 percent of study women said that they were not allowed to go for ANC check-ups. As respondents from Ramabai nagar slum area, stay on hill and therefore this may be the reason for not availing ANC services.

Antenatal care is essential for ensuring safe mother-




Figure 6. Reason for not receiving an antenatal check-up according to residence, Ramabai Nagar slum area, Bhandup.

Table 1. Utilization of antenatal care services showing antenatal care indicators in Ramabai Nagar Slum Hill, Bhandup, Mumbai.

Antenatal care indicators	Ramabai Nagar Slum, Bhandup	
	Total cases	Percent of cases
Percentage that received at least one antenatal check-up	310	89.6
Percentage that received three or more antenatal check-up	179	57.7
Percentage that received two or more T.T. injections	200	67.1
Percentage given any iron and Folic acid tablets or syrup	274	88.4
Tablets received/purchased		
1-40	138	50.4
41-100	76	27.7
100+	33	12.0
Not received	60	21.9
Median number of check-ups (for those who received at least one antenatal check-up)	3	
No antenatal check-ups	36	10.4

hood. During antenatal period, women are likely to face health problems of reproductive nature and there will be a package of measures available for expectant mothers, which ensures safe motherhood. The study women who have given at least one live birth during three years prior to the survey in the study area were considered to analyze the differentials in pregnancy problems experienced by mothers and the extent of utilization of antenatal care services.

This result suggests the need to inform women and families about the availability of free antenatal services in government health centres and about the benefits of antenatal check-ups to help overcome traditional attitudes and other hurdles that prevent them from seeking antenatal care during pregnancies. In addition,

since nearly one-fourth of the reasons reported deal with problems of accessibility, quality, and cost of services, utilization of antenatal check-ups could also be increased by lowering direct and indirect costs, improving quality and making services more accessible.

Components of antenatal care indicators

Table 1 indicates that 90 percent of study women received at least one antenatal check-up, 58 percent received three or more antenatal check-ups, 67 percent received two or more T.T. injections and 88 percent have consumed some kind of iron and Folic acid tablets or syrup. Median number of check-ups (for those who received at least one

Table 2. Utilization of antenatal care services showing components of antenatal check-ups in Ramabai Nagar Slum Hill, Bhandup, Mumbai.

Components of antenatal check-ups	Ramabai Nagar Slum, Bhandup			
	Cases		Cases	
	Yes	Percent	No	Percent
Weight measured	291	93.9	19	6.1
Height measured	233	75.2	77	24.8
Blood pressure checkup	263	84.8	47	15.2
Blood test	286	92.3	24	7.7
Urine test	284	91.6	26	8.4
Abdomen examined	280	90.3	30	9.7
X-Ray	134	43.2	176	56.8
Sonogram/Ultras	279	90.0	31	10.0
Any other test	105	33.9	205	66.1

Table 3. Utilization of antenatal care services showing antenatal advice in Ramabai Nagar Slum Hill, Bhandup, Mumbai.

Antenatal advice	Cases	
	Yes	Percent
Special diet	136	43.9
Danger sign for pregnancy	91	29.4
Delivery care	161	51.9
New born care	161	51.9
Family planning	136	43.9
Use of any form of tobacco	73	23.5
Walking exercise	131	42.3
Number of births for which the mother received at least one antenatal checkup	414	

antenatal check-up) is 3, and 10 percent of study women did not go for any antenatal check-up.

Components of antenatal check-ups

Antenatal measurements / Tests

Data on various components of antenatal check-ups undergone by women in Table 2 shows positive behavior pattern of women in utilizing antenatal care services in the study area. Weight measured is about 94 percent, blood and urine test is 92 percent, Sonogram and abdomen examination is 90 percent and blood pressure check-up is 85 percent.

Antenatal advice

Data on antenatal care advice in Table 3 reveals that the proportion of pregnant women in the study area who have not utilized proper advice on delivery care since danger sign for pregnancy is 29 percent, new born care is just 50 percent followed by special diet and family planning

which is about 44 percent. About 24 percent of women use tobacco during pregnancy.

Thus from the tables, it can be summed up that still, ten percent of women do not avail antenatal check-ups. Figure 5, Table 1 and 2 show quite a good MCH service performance. Data on antenatal care advice in Table 3 reveals that the proportion of pregnant women in the study area have not utilized proper advice on delivery care as new born care is just 50 percent followed by special diet and family planning is about 44 percent. Hence the study reveals excellent antenatal care seeking behavior of women towards antenatal measurements but moderate towards antenatal advice.

SOCIO-ECONOMIC CORRELATES

Table 4 shows the influence of the socio-economic characteristics of study women on the utilization of antenatal care services in the study area. It is clearly seen from the table that the utilization of antenatal care services increases with rise in the position of women with respect to each of the socio-economic factors. The utilization of

Table 4. Percentage that received antenatal care by selected background characteristics in Ramabai nagar slum, Bhandup, Mumbai.

Background characteristics	Ramabai Nagar Slum, Bhandup		
	Received antenatal checkup		Number of births
	Yes	Percent	
Mother's age			
15-24	130	91.5	175
25-29	119	89.5	160
30-49	61	85.9	79
Total			414
Birth order			
1	100	91.7	107
2	127	94.1	176
3	58	87.9	87
4+	25	69.4	44
Mother's education			
Illiterate	53	69.7	89
Literate, <middle school complete(1-6)	59	92.2	78
Middle school complete (7-9)	106	95.5	132
High school complete and above (10+)	92	96.8	115
Religion			
Hindu	211	90.2	277
Buddhist	78	91.8	107
Other	21	77.8	30
Caste			
SC and ST	108	87.8	150
OBC	43	84.3	63
Other	159	92.4	201
Standard of living			
Low	55	76.4	87
Medium	219	92.4	283
High	36	97.3	44
Total	310	-	414

antenatal care services decreases with rise in the birth order of children and rise in mother's age whereas, it increases with improvement in the position of women with respect to education and standard of living. It shows no fixed pattern in religion and caste. In fact the utilization of antenatal care services is almost same amongst Buddhist and Hindu and is quite high, about 90 percent. Whereas the utilization of these services is found to be low among OBC category of women compared with 'SC and ST' and 'Others' caste category.

In order to see the net effect of independent variables (description of the variables is provided in Table 5 for Ramabai nagar slum area) on the dependent variable which is dichotomous, logistic regression technique has been adopted.

DETERMINANTS OF UTILIZATION OF HEALTH CARE SERVICES DURING DELIVERY PERIOD (DELIVERY COMPLICATIONS): A LOGISTIC REGRESSION ANALYSIS

Logistic regression

Table 6 shows the influence of the reproductive health problems during pregnancy of study women who did not go for ANC on delivery complications in the study area. It is clearly seen from the table that delivery complications due to the pregnancy problems such as night blindness, anemia, any vaginal bleeding was 73.7 percent, whereas due to excessive bleeding it had shown 75% which was maximum, followed by blurred vision (64.7%),

Table 5. Measurement of variables used in the logistic regression analysis for Ramabai nagar slum area, Bhandup.

Category	Variables	Code
	Dependent variables	
Delivery complications	No (Ref)	0
	Yes	1
	Independent variables	
1. Night blindness	No (Ref)	0
	Yes	1
2. Blurred vision	No (Ref)	0
	Yes	1
3. Convulsions Not from fever	No (Ref)	0
	Yes	1
4. Swelling of the legs, body or face	No (Ref)	0
	Yes	1
5. Excessive fatigue	No (Ref)	0
	Yes	1
6. Anemia	No (Ref)	0
	Yes	1
7. Any vaginal bleeding	No (Ref)	0
	Yes	1
8. White discharge	No (Ref)	0
	Yes	1
9. Abdomen pain	No (Ref)	0
	Yes	1
10. Excessive bleeding	No (Ref)	0
	Yes	1
11. Excessive omitting	No (Ref)	0
	Yes	1
Utilisation of antenatal care service (ANC)	Antenatal Care service not utilized (Ref)	0
	Antenatal Care service utilized	1

convulsions not from fever (62.5%) and for the rest it was around 55%.

The logistic regression results in Table 7 reveals that the study women who have not gone for antenatal care and having pregnancy problems develop complications during the delivery. Here the dependent variable is complications at the time of delivery and the independent variables are problems during the pregnancy and study women who did not go for ANC. The results of logistic regression showed that excessive bleeding during pregnancy will influence complications during the delivery of the respondents from Ramabai nagar area.

CONCLUSION AND POLICY IMPLICATIONS

Maternal deaths are clustered around labour, delivery and the immediate postpartum period, with obstetric

haemorrhage being the main medical cause of death. Skilled attendance during delivery, access to emergency obstetric care and postnatal care (PNC) are cost effective and life saving investments for mothers. The extent of services available and availed during complications related to pregnancy, delivery and postpartum indicates the state of obstetric morbidity and mortality (Ministry of Health and Family Welfare, 2008).

This data shows that the extent of utilization of services pertaining to antenatal period is excellent for those other than illiterate women, low category SLI women and SC-ST, OBC women. The roll of socio-economic factors in service utilization is clearly evident in study area. It clearly shows that as education level increases, the utilization of ANC also increases. This study also reveals excellent antenatal care seeking behaviour of women towards antenatal measurements but moderate towards antenatal advice. About 35 percent of women from low

Table 6. Percentage reproductive health problems during pregnancy and ANC on delivery complications in Ramabai nagar slum, Bhandup, Mumbai.

Problems during pregnancy	Delivery complications	
	Yes	Percent
Night blindness	14	73.7
Blurred vision	22	64.7
Convulsions Not from fever	15	62.5
Swelling of the legs, body or face	60	55.6
Excessive fatigue	85	51.5
Anemia	11	73.3
Any vaginal bleeding	22	73.3
White discharge	29	55.8
Abdomen pain	53	57.0
Excessive bleeding	36	75.0
Excessive omitting	75	52.1
ANC	136	43.9

Table 7. Odds ratios from logistic regression examining the effect of antenatal care and reproductive health problems during pregnancy on delivery complications in Ramabai nagar slum, Bhandup, Mumbai.

Problems during pregnancy	Sig.	Odds ratio
Night blindness	0.405	1.800
Blurred vision	0.530	1.408
Convulsions Not from fever	0.197	1.405
Swelling of the legs, body or face	0.905	1.032
Excessive fatigue	0.401	1.714
Anemia	0.889	.951
Any vaginal bleeding	0.893	1.076
White discharge	0.181	1.927
Abdomen pain	0.694	1.121
Excessive bleeding	0.003	3.081**
Excessive omitting	0.373	1.239
ANC	0.744	0.873
Constant	0.201	0.563
-2 Log likelihood	442.771	
Cox and Snell R ²	0.094	
Nagelkerke R ²	0.126	
Number of births	414	

***p < 0.01, **P < 0.05.

category of standard of living are not availing delivery care services. This clearly indicates that there is a concentration of women amongst the poorest of economic stratum that goes without adequate maternal care.

Similarly the influence of the reproductive health problems during pregnancy having no ANC creates complications, while delivery and the most influencing

factor found was excessive bleeding.

Moreover, being on a hill, this slum area has no road infrastructure and public transport services. Further, hiring private transport is unaffordable for many slum-dwellers. The problem with transport is even worse if the referral is made at night because the cost challenge is compounded by the insecurity in slum communities.

Though community health volunteers (CHV) are playing a big role in communicating importance of ANC, delivery care and post-natal care to the expectant women, illiteracy and poverty prevent such women from utilizing health facilities.

Thus this paper suggests that there should be:

1. Effective awareness campaign through urban health centers.
2. Committed health workers.
3. Easy access to services.
4. Awareness about appropriate birth interval and proper diet during pregnancy.
5. Better health care delivery system.
6. Quality health care.
7. Follow-up care.

All these are for the betterment of reproductive health of women in such hilly slums particularly to the illiterates.

ACKNOWLEDGEMENTS

The author is thankful to his guide Dr. M. B. Joshi, Associate Professor, Government Medical College, Aurangabad. He is also grateful to Dr. C. P. Prakasam (retired.), Professor, International Institute for Population Sciences, Govandi and Mumbai for giving valuable suggestions while drafting this paper. He is also thankful to the funding agency, Ratan Tata Trust, Mumbai without whose help, this presentation would not have been possible. Above all the author is grateful to the conference organizers of the British Society for Population Studies, Professor Paul Boyle, Ms. Anne and Dr. Sabu Padmadas who have invited him to present this paper.

This paper was presented in the British Society for Population Studies (BSPS) Annual conference 2009 held from 9 - 11 September, 2009 at the University of Sussex, Falmer, Brighton, UK, organized by the British Society for Population Studies, based at the London School of Economics, London, UK.

Note:

a. Mumbai was selected for the trial, in view of presence of large urban slum population (highest among cities in Maharashtra). Situation analysis of the slums revealed existence of unlisted slums, pockets of underserved slum population and underutilization of existing health services.

b. The indexed women were the one who had a live birth during the last 3 years preceding the survey.

c. Based on this survey, the information was recorded as the self reported symptoms. This did not necessarily imply check up being carried out at the time of survey. d. Complete Antenatal Care for a pregnant mother includes

at least 3 antenatal check-ups, 2 doses of TT injection and consumption of 90+ IFA tablets prior to the expected date of delivery.

e. An antenatal check-up includes weight and height measurement, blood pressure, abdominal examination and diagnostic tests including urine test etc.

REFERENCES

- Anderson N (1960). *The Urban Community*, pp 191, Urban Land Policies, New York, United Nations, April 1952.
- Bang RA, Bang AT, Baitule M, Choudhary Y, Sarmikaddam S, Tale O (1989). High Prevalence of Gynaecological Diseases in Rural Indian women. *Lancet*. 1: 85-87.
- Bergel EE (1955). *Urban Sociology*, pp 410.
- Bhatia JC, Cleland J (1995a). Determinants of use of maternal care in a region of south India. *Health Trans. Rev.* 5(2): 127-142.
- Bhatia JC, Cleland J (1995b). Self reported symptoms of gynecological morbidity and their treatment in South India. *Stud. Family Plan.* 26(4):203-216.
- Bhatia JC, Cleland J (1996). Obstetric morbidity in South India, Results from a community survey. *Soc. Sci. Med.* 43 (10): 1507-1516.
- Carine R, Wendy JG (2006). Maternal mortality: who, when, where, and why. *Maternal Survival Series steering group. Lancet.* 368: 1189-200.
- Census of India (2001). Maharashtra population data with data on slum population in urban units.
- Directory of Slums (a) Slums came in into existence prior to year 1976 in Greater Mumbai. (b) Slums came in into existence between years 1976 to 1980 in Greater Mumbai. Published by the office of the additional collector (ENC), Mumbai & Mumbai Sub. Dist.
- Fronczak N, Arifeen SE, Moran AC, Caulfield LE, Baqui AH (2007). Delivery Practices of Traditional Birth Attendants in Dhaka Slums, Bangladesh. *J. Health. Populat. Nutr. Dec;* 25(4):479-487.
- Godbole VT, Talwalkar MA (1999). Programme for Children: An Assessment in Urban Areas of Maharashtra 1998. State Family Welfare Bureau, Pune.
- Harikrishnan KS (2009). High morbidity among women in Indian state. *One World South Asia.* 30 July 2009.
- Harrison KA (1990). The Political Challenge of Maternal Mortality in the Third World. *Maternal Mortality and Morbidity – A Call to women for action. Special Issue, May 28, 1990.*
- Jean CF, Alex E, Rose O (2008). Provision and Use of Maternal Health Services among Urban Poor Women in Kenya: What Do We Know and What Can We Do? *J. Urban Health.* 85(3): 428-442. 2008 May.
- Kapadia K, Tupe NR (2001). Do Women's Gender Attitudes Influence their Health? Evidence from Maharashtra, India. Paper under publication.
- Kapadia KN, Kanitkar T (2002). Primary Healthcare in Urban Slums. *EPW Commentary.* December 21, 2002.
- Khilare K (2001). Healthcare Services for Urban Population in Pimpri-Chinchwad Municipal Corporation, Unpublished paper.
- Koblinsky MA, Campbell OMR, Harlow D (1993). *Mother and More: A broader Perspective on Women's Health.* Koblinsky MA, Timyan J, Gay J (eds.). The Health of Women: A Global Perspective. Oxford: West View Press.
- Mayank S, Bahl R, Bhandari N (2001). Reproductive Tract Infections in Pregnant Women in Delhi, India. *Int. J. Gynecol. Obstet.* 75(1): 81-82.
- Ministry of Health and Family Welfare (1997); (1998b).
- Ministry of Health and Family Welfare. *Reproductive and Child Health – II programme.* New Delhi: Government of India, 2008.
- Montgomery, M. R. (2000) "Measuring Living Standards with Proxy Variables", *Demography*, vol. 37, no. 2, pp. 155-174
- Murthy N, Barua A (2001). Non-medical Determinants of Maternal Death in India. *Health Matters.* 9(17): 53-62. May 2001.
- Nangia S, Thorat S (2000): "Slums in a Metropolis, The Living Environment". Shipra Publications, Delhi NFHS-2(1998-99): International Institute for Population Sciences, Deonar, Mumbai.

- Parikh I, Taskar V, Dharap (1996). Gynaecological Morbidity among Women in a Bombay Slum. *Streehitakarini*. A Working Paper. pp 1-26. SNTD, Bombay.
- Prakasam CP (2004). Reproductive Morbidity among Adolescent Women in Andhra Pradesh and Tamil Nadu: Evidences from NFHS-2 data. *Women Health and Development*, Department of Population Studies, UGC-SAP (Phase-I), Sri Venkateswara University, Tirupati.
- Registrar General of India in Collaboration with Centre for Global Health Research, Canada. *Maternal mortality in India: 1997 – 2003, trends, causes and risk factors*. Sample Registration System, Registrar General of India and Centre for Global Health Research, University of Toronto, Canada, 2006.
- Registrar-General of India, 2007.
- Srinivasa DK, Narayana KA, Oumachigui A, Gautam R (1997). *Prevalence of Maternal Morbidity in a South Indian Community*. Unpublished Report. Pondicherry. JIPMER.
- United Nations Educational, Social and Cultural Organization (UNESCO). (*World Education Report, 1997*. Paris: UNESCO, 1997.
- UNICEF. *Maternal mortality – A woman dies every 5 minutes from child birth in India*. http://www.unicef.org/india/health_1341.htm (accessed Mar 3, 2009).
- Urban Female Sample Survey (1998a). Institute of Health Management (IHMP), Pachod, Pune Centre.
- World Health Organization (1996b).

Appendix

The standard of living is calculated by adding the following scores:

Morbidity caused by delivery location ($p = 0.58$).

Type of House: 4 for pucca, 2 for semi-pucca, 0 for kachha.

Toilet facility: 4 for own flush, 2 for public, 1 for public pit or open, 0 for no facility.

Source of lighting: 2 for electricity, 1 for other, 0 for no facility.