

Full Length Research Paper

# Molecular characterization of fowl adenoviruses associated with hydro-pericardium syndrome in broilers

Shah, M. S.<sup>1,2\*</sup>, Ashraf, A.<sup>3</sup>, Khan, M. I.<sup>1\*</sup>, Rahman, M.<sup>2</sup>, Habib, M.<sup>4</sup>, Babapoor, S.<sup>1</sup>, Ghaffar, A.<sup>2</sup>, Malik, I. R.<sup>2</sup>, Khannum, S. A.<sup>4</sup> and Qureshi, J. A.<sup>2,5</sup>

<sup>1</sup>Department of Pathobiology and Veterinary Sciences, University of Connecticut, Storrs, USA.

<sup>2</sup>Health Biotechnology Division, National Institute of Biotechnology and Genetic Engineering (NIBGE), Faisalabad, Pakistan.

<sup>3</sup>Department of Bioinformatics and Biotechnology, G.C University, Faisalabad, Pakistan.

<sup>4</sup>Animal Sciences Division, Nuclear Institute for Agriculture and Biology (NIAB), Faisalabad, Pakistan.

<sup>5</sup>Department of Clinical Biochemistry, College of Medicine, King Khalid University, ABHA, Kingdom of Saudi-Arabia.

Accepted 16 November, 2016

Hydropericardium syndrome (HPS) is a viral disease of poultry, which is common in hot and humid season during summer in Pakistan. Fowl adenovirus serotype 4 (FAdV-4) classified into the species Fowl adenovirus C and the genus Aviadenovirus, which were identified as the causative agent of the disease during field outbreaks. The virus was obtained from suspected broiler birds showing typical signs and postmortem lesions that is ascites, hydropericardium, multifocal hepatic necrosis and hepatitis. The virions were partially purified from the livers of infected birds, by sucrose density gradient ultracentrifugation. A variable part of the hexon gene (L1 and part of P1, 730 bp) was amplified by polymerase chain reaction (PCR) and its nucleotide sequence was determined. During nucleotide sequence analysis, the viruses were identified as FAdV- 4. The presence of FAdV-4 was also supported by scanning electron microscopy. Phylogenetic analysis of the partial hexon gene of the viruses determined their close relation with previously sequenced FAdV-4 isolates from India.

**Key words:** Fowl adenovirus serotype 4 (FAdV-4), hydro-pericardium syndrome, hexon, electron microscopy, phylogenetic analysis, taxonomy.

## INTRODUCTION

Hydro-pericardium syndrome (HPS) is an important disease of poultry particularly effecting 3 to 6 week old broiler chicks. The disease is characterized by sudden onset with 20 to 70% mortality, accumulation of fluid in the pericardial sac, enlarged discolored liver with necrotic foci and intra-nuclear inclusion bodies in hepatocytes (Dahiya et al., 2002; Anjum et al., 1989). The causative agent is fowl adenovirus serotype 4 (FAV-4), which

belongs to the genus Aviadenovirus of the family Adenoviridae. Adenoviruses are non-enveloped, medium sized, icosahedral particles with linear, double stranded DNA genome. Adenoviruses are classified into four recognized genera which include Atadenovirus, Siadenovirus, Mastadenovirus and Aviadenovirus while the only confirmed Fish adenovirus felled into fifth clade (Davison et al., 2003). Harrach et al. (2011) suggested to place the Turkey adenovirus species in genus Aviadenovirus and renamed the species in the family Adenoviridae by adding letter designation “*Possum adenovirus A*, *Goose adenovirus A*, *Canine adenovirus A*, *Tree shree adenovirus A* and *Frog adenovirus A*”. Transmission of the disease occurs through oral - fecal

\*Corresponding author. E-mail: [msalahuddin786@hotmail.com](mailto:msalahuddin786@hotmail.com). [mazhar.khan@uconn.edu](mailto:mazhar.khan@uconn.edu). Tel: +92 322 7748074. Fax: +92 41 2654213.

route, both vertically and horizontally. The first outbreak of HPS was reported in 1987 from *Angara Goth* near Karachi, Pakistan, so the disease was locally named as *Angara* disease (Anonymous, 1988; Roy et al., 2004). Later the syndrome was also found in medium weight layers and broiler breeders (Anjum, 1988). Afzal et al. (1991) and Kim et al. (2008) noticed that HPS was common in broilers at 3 to 5 week of age, whereas outbreaks were also reported in layers and breeders. The syndrome can be reproduced in broilers by the inoculation of a liver homogenate from affected birds (Anjum, 1990; Nakamura et al., 1999). Polymerase chain reaction (PCR) amplification of a hyper variable region (L1 and part of P1) of the hexon gene and its phylogenetic analysis has revealed that FAdV-4 isolates from chickens with HPS in Japan were identical but they were differentiated from the isolates of India and Pakistan (Meulemans et al., 2004; Masaji et al., 2009). This work was carried out to characterize the viruses causing HPS in Pakistan and to determine their phylogenetic relationship with other regional isolates.

## **MATERIALS AND METHODS**

### **Collection of infected samples**

Postmortems of suspected birds were conducted and livers were collected from naturally infected broilers showing typical post mortem lesions (enlarged liver, serous fluid in the pericardial sac and abdomen) of HPS. Details of samples are given in Table 1. One gram of liver was homogenized in 10 ml of phosphate buffer saline (PBS) and centrifuged at 3000 rpm for 15 min at 4°C. The supernatant was collected and stored at -20°C after addition of antibiotics that is Penicillin 10, 0000 I.U/ml, Streptomycin 10 mg/ml and Gentamycin 250 µg/ml (Roy et al., 2004).

### **Transmission of infection**

Ten healthy broiler birds of 3 weeks old from a commercial poultry farm were experimentally infected with 1 ml of 10 % liver homogenate and observed for signs and symptoms of the disease up to 10 days post inoculation, as shown in Table 2. Five broiler birds of same age were inoculated with 1 ml of sterile PBS and kept as control (Kumar et al., 1997). Birds were observed daily for clinical signs and mortality. At post mortem examinations, the livers were collected for re-isolation of the virus as described previously.

### **Semi-purification of the virus**

Partial purification of the virus was performed by ultra centrifugation of the supernatant, described in previous (Kumar and Chandra 2004). The supernatants were layered on preformed, 15 to 35% sucrose gradients prepared in 0.1 M PBS and centrifuged at 80,000 rpm for 2 h at 4°C. An opaque band was found between 25 and 35% sucrose gradient. This band was collected, diluted with PBS and spun again in a centrifuge at 80,000 rpm for 2 h at 4°C. The pellet was resuspended in 0.5 ml of PBS and stored at -20°C. Purity of the viral suspension was determined using spectrophotometer, by measuring the absorbance ratio at 260 and 280 nm as described by Kumar and Chandra (2004). This virus suspension was used for

DNA extraction.

### **Viral DNA extraction**

DNA was extracted as described by Ganesh et al. (2002) with some modifications. The semi purified viral suspension was digested by adding 200 µl lysis buffer (Tris-HCl, NaCl, SDS, and Proteinase K to final conc. of 25, 100 mM, 0.5% and 25 µg/ml respectively) in 200 µl viral sample. Subsequently, 100 µl of phenol:chloroform (a:a) was added and well mixed. After centrifugation, the supernatant was moved into a new tube, and the concentration of NaCl was adjusted at 0.2 M. DNA was precipitated by chilled (-20°C) ethanol. The DNA was pelleted by centrifugation. The DNA pellet was dried, re-dissolved in 20 µl of double-distilled water and stored at -20°C.

### **Detection of viral DNA by polymerase chain reaction (PCR)**

Amplification of a 730 bp variable part of the hexon gene was done by PCR to confirm the presence of viral DNA. The primers (FAVHL 5'- GACATGGGGTCGACCTATTTTCGACAT-3' and reverse primer FAVHR 5'- AGTGATGACGGGACATCAT-3') as well as the conditions of the PCR (35 cycles consisting of 94°C for 60 s, 57°C for 60 s and 72°C for 120 s) as described by Ganesh et al. (2001). The PCR products were analyzed along with 1 Kb DNA ladder on 1.5% agarose gels containing 0.1% ethidium bromide. The electrophoresis was performed at 100 Volts for 30 to 40 min in 1 X TAE buffer. After electrophoresis, PCR products were visualized by placing the gel on ultra violet light transilluminator.

### **DNA sequencing of the polymerase chain reaction (PCR) product**

Nucleotide sequence of the PCR products were determined from both sides using forward and reverse primers, on an ABI 3100 capillary sequencer. The sequence data were searched in public access databases (GenBank, EMBL) with BLAST (Basic Local Alignment Search Tool) as described by Altschul et al. (1990).

### **Scanning electron microscopy (SEM)**

Surface structure of the virus was analyzed under field emission scanning electron microscope (FESEM) as described by Hong et al. (2005) with some modifications. Virus particles were adsorbed on the surface of carbon coated grids by placing a drop of the semi purified viral suspension on the grid and drained out extra fluid. Specimens were negatively stained by adding a drop of 2% uranyl acetate stain for few seconds and extra stain was drained. Specimens were air dried and observed under Scanning Electron Microscope (Model No: JEOL JFM 7500 F). Micrograph was taken under low-dose conditions at 3.0 KV and at magnification of 30,000 x.

### **Phylogenetic analysis**

Multiple sequence alignment and phylogenetic analysis were done on observed divergence basis by Multiple Sequence Alignment program of "DNAMAN, Lynnon Biosoft" (Lynnon Corporation, Quebec, Canada). Pair-wise comparison of nucleotide sequences of the variable part of the hexon gene was done. A rooted genetic dendrogram was constructed by neighbor-joining method and phylogenetic relationships of our isolate with other reported isolates were determined. All sequences used for multiple alignments were obtained from public access databases GenBank and EMBL,

**Table 1.** Case history of flocks with natural outbreaks used for virus isolation.

No.	Year	Site of collection	Type of birds	No. of birds	Age of birds (days)	Mortality (%)
1	2008	Punjab, Pakistan	Broilers	8000	35	25
2	2009	Punjab, Pakistan	Broilers	6000	26	32

**Table 2.** Mortality results after experimental infection with 10% liver homogenate of diseased birds.

	Number of birds	Age of broilers	Infection		Mortality (%)
			Dose	Route	
<b>Treatment</b>	10	21 days	1 ml	S/C	80 (8/10)
<b>Control</b>	5	21 days	1 ml (PBS)	S/C	0 (0/5)

**Table 3.** Details of the sequence data used for multiple sequence alignment.

No.	Accession No.	Virus type	Host animal	Country of origin
1	AF207659.1	BAdV-6	Cattle	Iowa, USA
2	AF339923.1	FAdV-9	Chicken	Belgium
3	AY581275.1	FAdV	Turkey	Izatnagar, India
4	DM380829.1	CELO Virus	Chicken	Germany
5	DQ149638.1	HAdV-45	Human	Vienna, Austria
6	DQ264728.1	FAdV-4	Chicken	Pakistan
7	EF442329.1	Psittacine AdV-1	Parrot	Germany
8	EF685395.1	Fowl adenovirus C	Chicken	Canada
9	EU847626.1	FAdV-4	Chicken	Haryana, India
10	FN394664.1	FAdV-4	Chicken	Hemachal perdesch, India
11	L80007.1	EAdV-2	Horse	Davis, CA, USA

details are shown in Table 3.

## RESULTS AND DISCUSSION

During experimental infection, eighty percent mortality was observed in the treatment group while control birds remained healthy. Results are shown in Table 2. Similar results were found by Kumar et al. (1997) who experimentally infected the birds with FAdV-4.


Presence of the virus was confirmed by PCR amplification of a 730 bp fragment from the hexon gene as shown in Figure 1. Nucleotide sequence of the PCR products from two isolates revealed identical sequence that was submitted to the GenBank with accession no. FR686931.1.

The BLAST revealed that the nucleotide sequence of our isolate has 99% identity with several Indian isolates (Haryana, Hemachal Perdesch, Izetnagar; accession no. EU847626.1, FN394664.1, AY581275.1) of FAdV-4, while 98% identity was found with a Pakistani FAdV-4 isolate (accession no. DQ264728.1). The identity values with a German psittacine adenovirus (accession no. EF442329.1) and FAdV-1 (CELO) (accession no.


DM380829.1) were found to be 65% and 70% respectively.

Multiple sequence alignment (MSA) of the 730 bp variable region of the hexon gene with the corresponding gene fragment from other adenovirus is presented in Figure 4. MSA revealed that this region of human adenovirus serotype 45 contains 847 bp, and thus is longer than the corresponding part of other mastadenoviruses (equine or bovine). Our results are in line with the findings of Harrach et al. (2011) who found that, adenoviruses which infect the birds are classified into three genera including Atadenovirus, Mastadenovirus and Aviadenovirus. It was also found that the L1 – P1 region of the hexon gene is hypervariable in the middle both; the ends are more conserved thus being suitable for PCR primer design.

The results of phylogenetic analysis showed close relation with the regional (Indian and Pakistani) FAdV-4 isolates as supported by the tree on Figure 3. So it was suggested that isolates of FAdV in India and Pakistan had a common ancestor, which can be differentiated from other FAdV isolates of Canada, Belgium and Germany. Similarly, Meulemans et al. (2004) and Masaji et al. (2009) suggested that FAdV-4 isolates in Japan were



**Figure 1.** Agarose gel electrophoresis. Lane 1, + ve control; Lane 2, - ve control; Lane 3 and 4, 730 bp PCR product; Lane 5, 1 Kb molecular weight marker.




**Figure 2.** Scanning Electron micrograph of intact FAdV-4 showing icosahedral shape capsids, indicated by long arrows. Hexons found in back ground from disrupted viral particles indicated by small arrows. The scale bar represents 100 nm.

identical but they were different from the isolates of India and Pakistan.

The protein concentration of the semi-purified virus

samples was estimated to be 0.85 to 0.90  $\mu\text{g}/\mu\text{l}$ ., The SEM photograph of intact virions (Figure 2) revealed typical icosahedral capsids of the virus. The diameter of

0.05



**Figure 3.** Phylogenetic tree based on a 730 bp hypervariable region of the hexon (FAdV- 4 NIAB 01) a rooted genetic distance dendrogram.

each capsid was estimated to 70 to 80 nm. Laver et al. (1971) also found capsids of same shape and size of an avian adenovirus (CELO). The background of the photograph has small white particles which likely represent the group of hexons from disrupted viral particles as

previously found by Hong et al. (2005).

In summary, molecular characterization of two FAdV isolates from Pakistan revealed that these isolates were identical to each other but they were considerably different from isolates of other countries like Belgium and

AF207659.1\_BAdV- L80007.1\_EAdV-2 DQ149638.1\_HAdV- EF442329.1\_Psitt DM380829.1\_CELO\_ EF685395.1\_Fowl\_ AY581275.1\_FAdV\_ AF339923.1\_FAdV- FN394664.1\_FAdV- EU847626.1\_FAdV- DQ264728.1\_FAdV- FAdV-4\_NIAB\_01 Consensus 85 85 85 43 85 85 85 85 85 85 85

GATTGGGAAGTCTTATTTTGATATAGAAGGATTTTATAGAC A TTTTCT T A A TGGAA A TGCT T TC 85  
GACATGGGAGAAGTACTTTTACATCCGCGGGACCTAGAC C ACCCAGC C G T CAGC G GCAG C CC 85  
GACATGGCCAGCCTTACTTTGACATCCGCGGGCTCTGGAC C TCCAGC C A C CTGC C AGCT C CA 85  
..... ACCATCC C A G TGGT T AGCT T TC 43  
GACATGGGGGCGACCTACTTCGACATAAAGGGTGTGCTGGAC C ACCTTCC C G G CGGC A GGCT T TC 85  
GACATGGGGTTCGACCTATTTTCGACATCAAGGGAATCCTAGAC A GCCGTCC C G C CTGC C GGCT C CC 85  
GACATGGGGTTCGACCTATTTTCGACATCAAGGGAATCCTAGAC A GCCGTCC C G C CTGC C GGCT C CC 85  
CA GG GTCG. TTC GAC AT( GGAAT CT AGT CC T T T A T GG C CC 85  
..... T A T T 85  
..... T 85  
..... T 85  
GACATGGGGTTCGACCTATTTTCGACATCAAGGGAATCCTAGAC A GCCGTCC C G C CTGC C GGCT C CC 85  
Consensus cg gg tt aa cc ta gg ac ta aa

AF207659.1\_BAdV- L80007.1\_EAdV-2 DQ149638.1\_HAdV- EF442329.1\_Psitt DM380829.1\_CELO\_ EF685395.1\_Fowl\_ AY581275.1\_FAdV\_ AF339923.1\_FAdV- FN394664.1\_FAdV- EU847626.1\_FAdV- DQ264728.1\_FAdV- FAdV-4\_NIAB\_01 Consensus 139 140 170 94 139 142 142 142 142 142 142

CCC C T T T AATCAG TATGCCAATATAGGCTTTTC ..... AATAACGAGGATA 139  
AAT G T T C AGGGCG GCCAACCCATGTCTAGTA..... CCA ..... AAAATAACTACTAC 140  
GCC G C C AGAGCG TCCCAATCCAGCCAGTGGGATGCAAAGGAAAAGGAAGGAGTTGCCCAA CAGAAAAAATGTTTT 170  
CTC G T T AGGAAG CATCTTTAACAGTTGGGCCG..... T..... GCGGACAA 94  
CCC T G A GAGAAG TATTTCAACACCTGGGTG..... A..... GCACT.GGTCTC 139  
CGC G T C AGGAGT CATGTTTAAACAACCTGGTCCG..... A..... C.GGCGCCCGGGCAGA 142  
CGC G T C AGGAGT CATGTTTAAACAACCTGGTCCG..... A..... C.GGCACCCGGGCAGA 142  
GG GT TT AC TC ..... A..... G C C C G A A 142  
G T T A C G ..... A..... G C C G A A 142  
G T T A C G ..... A..... G C C C G A A 142  
CGC G T C AGGAGT CATGTTTAAACAACCTGGTCCG..... A..... C.GGCACCCGGGCAGA 142  
CGC G T C AGGAGT CATGTTTAAACAACCTGGTCCG..... A..... C.GGCACCCGGGCAGA 142  
Consensus t gc cc a c a

AF207659.1\_BAdV- L80007.1\_EAdV-2 DQ149638.1\_HAdV- EF442329.1\_Psitt DM380829.1\_CELO\_ EF685395.1\_Fowl\_ AY581275.1\_FAdV\_ AF339923.1\_FAdV- FN394664.1\_FAdV- EU847626.1\_FAdV- DQ264728.1\_FAdV- FAdV-4\_NIAB\_01 Consensus 212 213 255 167 212 212 212 212 212 212 212

CAGTTTATTTAGCTCAATTCCTCAAATTTA CTGCTAATGAAAAGGAGTGAACAAGCTATGGCTATAAA..... 212  
AGTGACCTTTGGCCAAGCG CATTTCGAAGTTCGAGATACCCGCTAGCGGGCTGCGCATCGCCCTCGACAAA..... 213  
AAAAACATTTGGTGTGGCC CTACAGGTGGTT TAATATTACAGATCAGGGTTTGTACTTGGAACTGAGGAAACAGCTGAAAAC 255  
ACACTACCGTAGTCGGGCA ATGCCAATGTGC ACACGAACGACACAACACCACCCCGGTGCCACCGCTAC..... 167  
AGACCAATGTGGTGGGACA ATGACCAACGTGC ACACAAATCAGACCAGGAACGACAAGACGGCCACGCTTCA..... 212  
ACGTGTCCGCTCCGGTCA CTGTCCAATGTGC ATACCAACACGACACCACCAAAGACAACGACGGCGGC..... 212  
ACGTGTCCGCTCCGGTCA CTGTCCAATGTGC ATACCAACACGACACCCTCCAAGACACGACGGCGGC..... 212  
GT A CT CG A TGC CC TAC CA CACGAAC CC AAG AC AC GC ..... 212  
T T T G H G X A A A G A C C AG C C C ..... 212  
T T T G H G X A A A G A C C AG C C C ..... 212  
ACGTGTCCGCTCCGGTCA CTGTCCAATGTGC ATACCAACACGACACCCTCCAAGACACGACGGCGGC..... 212  
ACGTGTCCGCTCCGGTCA CTGTCCAACGTGC ATACCAACACGACACCCTCCAAGACACGACGGCGGC..... 212  
Consensus g t

AF207659.1\_BAdV- L80007.1\_EAdV-2 DQ149638.1\_HAdV- EF442329.1\_Psitt DM380829.1\_CELO\_ EF685395.1\_Fowl\_ AY581275.1\_FAdV\_ AF339923.1\_FAdV- FN394664.1\_FAdV- EU847626.1\_FAdV- DQ264728.1\_FAdV- FAdV-4\_NIAB\_01 Consensus 278 270 340 233 280 279 279 279 279 279 279

.....TGTAAGTCTACTACTCTTAATCCACAGACTGGTAGGA..CAGATGATATTCTGGACCAATGATAT.. 278  
.....TACTGCGACCAACTTACAGCCAGAACCCTCAGATTGG...AATCGT TCGTGGGA.....GCC... 270  
GTTAAAAAGGATATCTATGACGAGAAACCTTACAGCCTGAACCTCAAGTTGGTGAAGAAA TGGCAGSAAAGTGAAGCCTTTT 340  
.....CATCATGCGCAACTTTT TGGCATAAATCCCAATGTCAATTGAGGAC CGGCAT.ATCAGAATACAC... 233  
.....GCAGGTCAATAGCATCT CGGGGTGGTTCCCAACGTCAACCTGGGAC CGGCCTCAGTCAACTAGCAT.. 280  
.....GCAGGTGACGAAGATTT CGGCCTTTCCCAACCCCAACCAGGGA CGGAATAAATCCTCTGCGG... 279  
.....GCAGGTGACGAAGATTT CGGCCTTTCCCAATCCCAACCAGGGA CGGAAGAAATCCTCTGCGA... 279  
.....GCAGGTGACGAAGATTT CGGCCTTTCCCAATCCCAACCAGGGA CGGAGTAAATCCTCTGCGG... 279  
.....CAGG GA A A ATTT G C T TCC CAI CC CAI CCAG GA C GA CC CT CGA... 279  
..... .A... 279  
..... .A... 279  
..... .A... 279  
..... .A... 279  
Consensus c c

AF207659.1\_BAdV- L80007.1\_EAdV-2 DQ149638.1\_HAdV- EF442329.1\_Psitt DM380829.1\_CELO\_ EF685395.1\_Fowl\_ AY581275.1\_FAdV\_ AF339923.1\_FAdV- FN394664.1\_FAdV- EU847626.1\_FAdV- DQ264728.1\_FAdV- FAdV-4\_NIAB\_01 Consensus 348 345 425 309 354 342 342 342 342 342 342

.....TGATTCTGCAAAAAATGGAGGCTATGGGCGAATTCTTAGTCTGAAA.....GCAAGGGTGTAAATTTCCAGCT 348  
.....GGAACGACGCTTAGCGACGACAT.....TGCCCGCTGGGAC GCGCTTCTTAAACAA AGCAGAACCCACCCTGTG 345  
ATGGAGGAAGGGCTATTAAGAAAGACCAAAAATGAAGCCATGCTAT GTTCATTGGCAGAC ACTAATGAAAAGGAGGACA 425  
.....CTAGCGGTAACACCACGTTAATGCCATAGGAGTATCC CCAAGTTCGC.ACAG GGCTGGTACTAACTTAG.CT 309  
.....C...CCGGCCGACGTGGATAATATGGCGTGGTGGGACGTTT CCAAGGTAG...AC AGCGGGCGTGAAGCAGG.CG 354  
.....CAGGTAGAAAACGCCAACACCCGGCGTCTCGGTCGCTTC CCAAGT.....CTAGTAC.....AATTACG.CT 342  
.....CAGGTAGAAAACGCCAACACCCGGCGTCTCGGTCGCTTC CCAAGT.....C GGTAC.....AATTACG.CT 342  
.....CAGGTAGAAAACGCCAACACCCGGCGTCTCGGTCGCTTC CCAAGT.....C GGTAC.....AATTACG.CT 342  
..... GG AC CAAC GCGG TCT GCT CC CA AGT..... TA G.CT 342  
..... T..... G T 342  
..... T..... G T 342  
..... T..... G T 342  
..... T..... G T 342  
Consensus g c




Figure 4. MSA of 730 bp variable region of Hexon gene from various Adenovirus isolates.

Canada. It was also determined that Pakistani isolates had nearest phylogenetic relationship with Indian.

## ACKNOWLEDGEMENT

This work was supported by the Higher Education Commission, Government of Pakistan.

## REFERENCES

- Afzal M, Muneer R, Stein G (1991). Studies on the aetiology of hydropericardium syndrome (Angara disease) in broilers. *Vet. Rec.*, 128: 591–593.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990). Basic local alignment search tool. *J. Mol. Biol.*, 215: 403 – 410.
- Anjum AD (1990). Experimental transmission of hydropericardium syndrome and protection against it in commercial broiler chickens. *Avian Pathol.*, 19: 655–660.
- Anjum AD, Sabri MA, Iqbal Z (1989). Hydropericarditis syndrome in broiler chickens in Pakistan. *Vet. Rec.*, 124: 247–248.
- Anjum AD (1988). Study of pathogenicity of the hydropericardium syndrome agent. Proceedings of National Seminar on Hydropericardium syndrome in chickens in Pakistan, Rawalpindi, pp. 111-115.
- Anonymous (1988). Proceedings of 1<sup>st</sup> National Seminar on Hydropericardium-pulmonary-edemacum-hepatonephritis-syndrome, Karachi; Pakistan Vet. Med. Assoc., pp. 1-100.
- Dahiya S, Srivastava RN, Hess M, Gulati BR (2002). Fowl adenovirus serotype 4 associated with outbreaks of infectious hydropericardium in Haryana, India. *Avian Dis.*, 46: 230–233.
- Davison AJ, Benko M, Harrach B (2003). Genetic content and evolution of adenoviruses. *J. Gen. Virol.*, 84(11): 2895- 2908.
- Ganesh K, Suryanarayana V, Raghavan R, Gowda S (2001). Nucleotide sequence of L1 and part of P1 of hexon gene of fowl adenovirus associated with hydropericardium syndrome differs with the corresponding region of other fowl adenoviruses. *Vet. Microbiol.*, 78: 1–11.
- Ganesh K, Suryanarayana V, Raghavan R (2002). Detection of Fowl adenovirus associated with Hydro-pericardium Syndrome by a Polymerase chain reaction. *Vet. Res. Commun.*, 26(1): 73-80.
- Ganesh K, Raghavan R, Gowada S, Satyanarayana ML, Suryanarayana VVS (2002). Purification and characterization of the aetiological agent of Hydropericardium Hepatitis Syndrome from infected liver tissue of broiler chickens. *Trop. Anim. Health Prod.*, 34: 7-17.
- Harrach B, Benkő M, Both GW, Brown M, Davison AJ, Echavarría M, Hess M, Jones MS, Kajon A, Lehmkuhl HD, Mautner V, Mittal SK, Wadell G (2011). Family *Adenoviridae*. King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (eds) *Virus Taxonomy: Classification and Nomenclature of Viruses*. 9<sup>th</sup> report of the International Committee on Taxonomy of Viruses. Elsevier, San Diego pp. 95-111.
- Hong SS, Szolajska E, Schoehn G, Franqueville L, Myhre S, Lindholm L, Ruigrok RW, Boulanger P, Chroboczek J (2005). The 100K-chaperone protein from Adenovirus serotype 2 (Subgroup C) assists in trimerization and nuclear localization of Hexons from Subgroups C and B Adenoviruses. *J. Mol. Biol.*, 352: 125 – 138.
- Kim NJ, Seong HB, Min JK, Jeom JK, Haan WS, InPil M (2008). Outbreaks of Hydropericardium Syndrome and Molecular Characterization of Korean Fowl Adenoviral Isolates. *Avian Dis.*, 52(3): 526-530.
- Kumar R, Chandara R (2004). Studies on structural and immunogenic polypeptides of hydropericardium syndrome virus by SDS-PAGE and western blotting. *Comp. Immun. Microbiol. Infect. Dis.*, 27: 155–161
- Kumar R, Chandara R, Shukla SK, Agrawal DK, Mahesh K (1997). Hydro-pericardium Syndrome (HPS) In India: A Preliminary study on the causative agent and control of the disease by inactivated autogenous vaccine. *Trop. Anim. Health Prod.*, 29: 158-164.
- Laver WG, Youngusband HB, Wrigley NG (1971). Purification and properties of chick embryo lethal orphan virus (an avian adenovirus). *Virology*, 45: 598–614.
- Masaji M, Masaki C, Toshikazu I, Kikuyasu N, Shigeo Y, Tadao I (2009). Genetic Characterization of Fowl Adenoviruses Isolated from Chickens with Hydropericardium Syndrome in Japan. *J. Vet. Med. Sci.*, 71(11): 1455–1458.
- Meulemans G, Couvreur B, Decaesstecker M, Boschmans M, vanden Berg TP (2004). Phylogenetic analysis of fowl adenoviruses. *Avian Pathol.*, 33:164–170.
- Nakamura K, Mase M, Yamaguchi S, Shibahara T, Yuasa N (1999). Pathologic study of specific-pathogen-free chicks and hens inoculated with adenovirus isolated from hydropericardium syndrome. *Avian Dis.*, 43: 414–423.
- Roy P, Vairamuthu S, Sakthivelan SM, Purushothaman V (2004). Hydropericardium syndrome in Japanese quail (*Coturnix coturnix japonica*). *Vet. Rec.*, 155: 273–274.
- Roy P, Muralimanohar B, Koteeswaran A, Omprakash AV (2004). Experimental studies on hydropericardium syndrome in two different synthetic lines of broiler chickens. *Vet. Archive*, 74(2): 157-164.