

## Full Length Research Paper

# Survey of plant-parasitic nematodes associated with yams in Edo, Ekiti and Oyo states of Nigeria

A. A. Adegbite<sup>1</sup>, J. O. Saka<sup>1</sup>, G. O. Agbaje<sup>1</sup>, O. F. Owolade<sup>1</sup>, G. O. Olaifa<sup>2</sup>, A. Lawal<sup>3</sup> and S. T. Ojo<sup>4</sup>

<sup>1</sup>Institute of Agricultural Research and Training, Obafemi Awolowo University, P. M. B. 5029, Moor-Plantation, Ibadan-Nigeria.

<sup>2</sup>Department of Plant Science, Obafemi Awolowo University, Ile-Ife.

<sup>3</sup>Oyo State Agricultural Development Programme, Zonal Headquarters, Oyo.

<sup>4</sup>Ekiti State Agricultural Development Programme, Ikole-Ekiti.

Accepted 19 May, 2019

A survey was conducted to determine the types, frequency and population of plant parasitic nematodes associated with the soils and roots of Yam (*Dioscorea* spp.) in all Local Government Areas of Edo, Ekiti and Oyo States of Nigeria using random sampling for soil and root and using pie pan modification of Baerman funnel for plant parasitic nematode extraction. Twelve, eleven and ten genera of plant parasitic nematodes were encountered in the three States respectively, while ten genera each were identified from root samples from the three States. Plant-parasitic nematodes recovered included *Meloidogyne* spp., *Pratylenchus* spp., *Scutellonema* spp., *Radopholus* spp., *Aphelenchoides* spp., *Trichodorus* spp., *Rotylenchus* spp., *Helicotylenchus* spp., *Aphelenchus* spp., *Longidorus* spp., *Xiphinema* spp and *Rotylenchulus* spp. *Pratylenchus* spp., *Scutellonema* spp., and *Meloidogyne* spp were most widely distributed with frequency rating of 70, 60 and 55% respectively in soil samples from Edo State and in the root samples the three genera predominated with 75, 60 and 60% frequency rating respectively. *Scutellonema* spp., *Meloidogyne* spp., and *Pratylenchus* spp were most widely distributed with a frequency rating of 75, 70 and 60% respectively in soil samples from Ekiti State and in the root samples the three genera predominated with 70, 65 and 50% frequency rating respectively while *Meloidogyne* spp., *Scutellonema* spp., and *Pratylenchus* spp were mostly widely distributed with a frequency rating of 70, 65 and 62.7% respectively in soil samples from Oyo State and in the root samples the three genera predominated with 65, 60 and 60% frequency rating respectively.

**Key words:** - Yam (*Dioscorea* spp.), types, frequency, population of plant parasitic nematodes, *Meloidogyne* spp., *Pratylenchus* spp., *Scutellonema* spp., *Radopholus* spp.

## INTRODUCTION

Yams (*Dioscorea* spp.) constitute one of the most important food crops in the tropics and most important group of staple foods especially in the yam zone (25°N and 25°S) of West Africa (Coursey, 1967). There are between 300 and 600 species of *Dioscorea*, of which only a few are edible. The four most commonly cultivated are the water yam (*D. alata* L.), yellow yam (*D. cayenensis* Lam.), Chinese yam (*D. esculenta* [Lour.] Burk.) and the

white yam (*D. rotundata* Poir). *D. cayenensis* and *D. rotundata* are indigenous to West Africa, *D. alata* and *D. esculenta* are native to Asia. *D. rotundata* is the most important species of yams in Africa; followed by *D. cayenensis*. The bulk of global yam production is concentrated in West Africa, with Nigeria producing the largest proportion followed by Ghana and Cote d'Ivoire (FAO-STAT 2004). Yam is the second most important root and tuber crop in the World after cassava in terms of production, and contributes more than 200 dietary calories per day for 60 million people in the yam zone (Nweke et al., 1991). In Nigeria, yam production is undertaken mainly in the South-eastern zone, the South-Western zone, the

\*Corresponding author. E-mail: [r.ayoadegbite@amuromail.com](mailto:r.ayoadegbite@amuromail.com), [talk2triple\\_a@fastmail.ca](mailto:talk2triple_a@fastmail.ca).

area south of River Benue and east of River Niger, which stretches from the humid forest to the subhumid guinea savannah ecological zones (Nweke et al., 1991) where the soil fertility and rainfall permit their production. Yams require rainfall ranging between 1168.4 and 2500 mm per annum. Yam productions are expanding annually but realized yields are usually considerably lower than potential yields.

Plant- parasitic nematodes damage is an important factor in tuber quality reduction and yield loss in yam in the field and in storage. A large number of plant- parasitic nematodes associated with yam cultivation have been reported from various yam producing areas of the world (Ayala and Acosta, 1971; Bridge, 1972; Thompson et al., 1973; Adesiyan and Odihirin, 1977; Caveness, 1982; Hahn et al., 1989; Green and Florini, 1996; Agbaje et al., 2002; 2003; Adegbite et al., 2005).

These are the yam nematode *Scutellonema bradys*, the root-knot nematode *Meloidogyne* spp. and the lesion nematode *Pratylenchus* spp., which are all field and post-harvest pests (Caveness, 1982; Hahn et al., 1989; Agbaje et al., 2002; 2003; Adegbite et al., 2005). However, plant parasitic nematodes associated with yam in South Western Nigeria, particularly in Edo, Ekiti and Oyo States have not been fully investigated. This survey was conducted to update available information on nematodes of yam in South Western Nigeria and determine the types, population levels and frequency of occurrence of plant parasitic nematodes associated with yam in the three states.

## MATERIALS AND METHODS

The survey was carried out in Edo, Ekiti and Oyo States in South Western Nigeria between January and June 2006. All the Local Government Areas of the three States were visited for collection of soil and root samples of yam which consists of 20, 18 and 33 Local Government Areas respectively. In each of the Local Government Areas, farms in three towns were randomly selected for sampling making a total of 60, 54 and 99 farms sampled in Edo, Ekiti and Oyo States respectively. In each farm, 10 plants were randomly selected for sampling. Soil samples from around the roots of each plant were collected using a soil auger along the four cardinal directions at the base of each plant in order to cover as much of the rhizosphere as possible. Soil samples were collected to a depth of 15-30cm with garden trowels. Samples (soil and root) from each farm were pooled and sealed in plastic bags and protected from the sun (Ricka and Barker, 1992). The samples were properly labeled and taken to Plant Protection Laboratory of the Institute of Agricultural Research and Training, Moor Plantation, Ibadan-Nigeria for analysis and for identification of plant parasitic nematodes.

Plant parasitic nematodes were extracted from the soil using the pie-pan modification of the Baerman Funnel method (Southey, 1986). Each composite soil samples was mixed thoroughly and plant parasitic nematodes were extracted from 200ml sub-sample. The set up was left undisturbed for 24 h before decanting the suspension into a beaker. Ten extraction trays were set up per sample. Plant parasitic nematodes in each suspension were killed by adding an equal volume of hot water to the nematodes suspension and each sample was then adjusted to a desired volume. The suspension was thoroughly mixed using a magnetic stirrer and 5ml aliquot was drawn from each suspension into a Doncaster

counting dish for identification and quantification of the extracted nematodes. Temporary mounts of nematodes were prepared prior to nematode identification. Identification of plant parasitic nematodes to the generic level was done using the Lucid-key of Bell (2004). In case of the root samples 10g/composite root samples were cut into pieces and macerated with an electric blender for 30seconds. Samples were then processed and identified in the same way as for the soil samples. Percentage frequency was determined using the formula  $n/N \times 100$ .

Where  $n$  = the number of times an individual nematode occurred in all the samples and  $N$  is the sample size (60 for Edo State, 54 for Ekiti State and 99 for Oyo State). Also the percentage nematode population was determined using  $In/TN \times 100\%$ .

Where  $In$  = the individual nematode population in all the samples while  $TN$  is the total population of all the nematodes extracted in all the samples.

## RESULTS

Twelve and ten genera of plant parasitic nematodes were encountered in soil and root samples collected from Edo State respectively. In the soil, plant parasitic nematodes identified were *Pratylenchus* spp., *Scutellonema* spp., *Meloidogyne* spp., *Aphelenchoides* spp., *Helicotylenchus* spp., *Rotylenchus* spp., *Radopholus* spp., *Trichodorus* spp., *Aphelenchus* spp., *Rotylenchulus* spp., *Longidorus* spp., and *Xiphinema* spp. *Pratylenchus* species was the most frequently occurring species in the soil (70%) where the population was 15500/200 ml soil which was followed by *Scutellonema* species with 60% frequency rating and a population of 12355/200 ml soil while *Meloidogyne* species had frequency rating of 57% and a population of 11542/200 ml soil. *Longidorus* and *Xiphinema* species had the lowest frequency rating of 16.5% while *Rotylenchulus* species had the lowest population of 1245/ 200 ml soil (Table 1). In the root sample similar trends were observed in which *Pratylenchus* spp was the most frequently encountered species (75%) and a population of 8500/10 g root, followed by *Scutellonema* and *Meloidogyne* species with 60% frequency ratings respectively (Table 2).

Eleven and ten genera of plant parasitic nematodes were encountered in soil and root samples collected from Ekiti State respectively. In the soil, plant parasitic nematodes identified were *Scutellonema* spp., *Meloidogyne* spp., *Pratylenchus* spp., *Trichodorus* spp., *Aphelenchus* spp., *Helicotylenchus* spp., *Radopholus* spp., *Longidorus* spp., *Xiphinema* spp., *Rotylenchulus* spp and *Aphelenchoides* spp. *Scutellonema* spp was the most frequently occurring species in the soil (76%) where the population was 19550/200 ml soil which was followed by *Meloidogyne* spp with 70% frequency rating and a population of 17655/200 ml soil while the third frequently occurring species was *Pratylenchus* with frequency rating of 63% and a population of 15887/200 ml soil. *Xiphinema* spp had the lowest frequency rating of 21% while *Aphelenchus* spp had the lowest population of 1845/200 ml soil (Table 3).

In the root, similar trends were observed in which *Scu-*

**Table 1.** Plant Parasitic Nematodes extracted from soils around the roots of Yam in Edo State

Nematode Genera	Frequency of Occurrence	% Frequency Rating*	Nematode Population/200 ml soil	% Nematode Population **
<i>Pratylenchus</i> spp	42	70	15500±175.6	18.2
<i>Scutellonema</i> spp	36	60	12355±163.5	14.5
<i>Meloidogyne</i> spp	34	57	11542±160.8	13.5
<i>Aphelenchoides</i> spp	30	50	8690±120.5	10.2
<i>Rotylenchus</i> spp	20	33	6538±100.2	7.7
<i>Radopholus</i> spp	18	30	6015±95.5	7.1
<i>Helicotylenchus</i> spp	15	25	5835±87.6	6.8
<i>Aphelenchus</i> spp	15	25	5835±87.6	6.8
<i>Trichodorus</i> spp	13	22	4567±71.6	5.3
<i>Xiphinema</i> spp	9	15	3695±73.8	4.3
<i>Longidorus</i> spp	9	15	3468±72.5	4.1
<i>Rotylenchulus</i> spp	6	10	1245±68.9	1.5

\*  $n/N \times 100$  (n = number of times individual nematodes occurred and N = Sample size (60)).

\*\*  $In/N \times 100$  (n = Individual nematode in all the samples and TN = Total Population of all the nematodes extracted in all the samples).

**Table 2.** Plant Parasitic Nematodes extracted from the roots of Yam in Edo State

Nematode Genera	Frequency of Occurrence	% Frequency Rating *	Nematode Population/10 g root	% Nematode Population **
<i>Pratylenchus</i> spp	45	75	8500±250.5	14.7
<i>Scutellonema</i> spp	36	60	7350±245.8	12.7
<i>Meloidogyne</i> spp	34	60	6990±238.5	12.1
<i>Rotylenchus</i> spp	16	27	6535±230.8	11.3
<i>Radopholus</i> spp	13	22	6285±200.5	10.8
<i>Helicotylenchus</i> spp	12	20	6158±198.8	10.6
<i>Aphelenchus</i> spp	10	17	5215±195.9	9.0
<i>Trichodorus</i> spp	10	17	4815±195.8	8.3
<i>Xiphinema</i> spp	8	13.3	3965±190.9	6.8
<i>Longidorus</i> spp	5	8.3	2156±185.6	3.7

\*  $n/N \times 100$  (n = number of times individual nematodes occurred and N = Sample size (60)).

\*\*  $In/N \times 100$  (n = Individual nematode in all the samples and TN = Total Population of all the nematodes extracted in all the samples).

*tellonema* spp was the most frequently encountered species (70%) and a population of 7450/10 g root, followed by *Meloidogyne* and *Pratylenchus* species with frequency ratings of 65 and 52% respectively (Table 4).

Ten genera of plant parasitic nematodes were encountered in soil and root samples collected from Oyo State respectively. In the soil, plant parasitic nematodes identified were *Meloidogyne* spp., *Scutellonema* spp., *Pratylenchus* spp., *Aphelenchoides* spp., *Trichodorus* spp., *Radopholus* spp., *Helicotylenchus* spp., *Rotylenchus* spp., *Longidorus* spp and *Xiphinema* spp. *Meloidogyne* spp was the most frequently occurring species in the soil (70%) with a population of 20565/ 200 ml soil which was followed *Scutellonema* spp with 65% frequency rating and a population of 17555/200 ml soil while the third frequently occurring species was *Pratylenchus* spp with frequency rating of 62.7% and a population of 16543/200

ml soil. *Longidorus* spp and *Xiphinema* spp had the lowest frequency rating of 15% with populations of 1234/200 ml soil respectively (Table 5). In the root, similar trends were observed in which *Meloidogyne* spp was the most frequently encountered species (65%) and a population of 8765/10 g root, followed by *Scutellonema* spp and *Pratylenchus* spp with frequency ratings of 60% respectively (Table 6).

## DISCUSSION

Yams are food crops of major importance in tropical agriculture and they provide the staple food stuff for millions of people in many tropical countries, most notably in West Africa, the Caribbean area and parts of South East Asia.

The intensification of agriculture has led to continuous

**Table 3.** Plant Parasitic Nematodes extracted from soils around the roots of Yam in Ekiti State

Nematode Genera	Frequency of Occurrence	% Frequency Rating*	Nematode Population/200 ml soil	% Nematode Population**
<i>Scutellonema</i> spp	41	76	19550±234	16.2
<i>Meloidogyne</i> spp	38	70	17655±217	14.6
<i>Pratylenchus</i> spp	34	63	15887±199	13.2
<i>Trichodorus</i> spp	20	37	8765±165	7.3
<i>Aphelenchus</i> spp	24	44	1845±56	1.5
<i>Helicotylenchus</i> spp	32	59	12345±189	10.1
<i>Radopholus</i> spp	15	28	10543±146	9.0
<i>Longidorus</i> spp	10	19	5467±89	4.5
<i>Xiphinema</i> spp	10	21	7654±100	6.4
<i>Rotylenchulus</i> spp	13	24	9876±125	8.2
<i>Aphelenchoides</i> spp	30	56	10897±155	9.0

\*  $n/N \times 100$  (n = number of times individual nematodes occurred and N = Sample size (54)).

\*\*  $In/N \times 100$  (n = Individual nematode in all the samples and TN = Total Population of all the nematodes extracted in all the samples).

**Table 4.** Plant Parasitic Nematodes extracted from the roots of Yam in Ekiti State.

Nematode Genera	Frequency of Occurrence	% Frequency Rating *	Nematode Population/10 g root	% Nematode Population **
<i>Scutellonema</i> spp	38	70	19550±234	16.2
<i>Meloidogyne</i> spp	35	65	17655±217	14.6
<i>Pratylenchus</i> spp	28	52	15887±199	13.2
<i>Trichodorus</i> spp	15	28	8765±165	7.3
<i>Aphelenchus</i> spp	13	24	1845±56	1.5
<i>Helicotylenchus</i> spp	12	22	12345±189	10.1
<i>Radopholus</i> spp	10	19	10543±146	9.0
<i>Longidorus</i> spp	5	9	5467±89	4.5
<i>Xiphinema</i> spp	5	9	7654±100	6.4
<i>Rotylenchulus</i> spp	13	24	9876±125	8.2

\*  $n/N \times 100$  (n = number of times individual nematodes occurred and N = Sample size (54)).

\*\*  $In/N \times 100$  (n = Individual nematode in all the samples and TN = Total Population of all the nematodes extracted in all the samples).

**Table 5.** Plant Parasitic Nematodes extracted from soils around the roots of Yam in Oyo State

Nematode Genera	Frequency of Occurrence	% Frequency Rating *	Nematode Population/200 ml soil	% Nematode Population **
<i>Meloidogyne</i> spp	69	70	20565±156	22
<i>Scutellonema</i> spp	64	65	1755±148	19
<i>Pratylenchus</i> spp	62	62.7	16543±130	18
<i>Aphelenchoides</i> spp	52	52.5	8958±66	10
<i>Trichodorus</i> spp	45	45.5	7335±50	8
<i>Radopholus</i> spp	40	40.4	7056±45	8
<i>Helicotylenchus</i> spp	35	35.4	6592±40	7
<i>Rotylenchus</i> spp	30	30.3	5996±40	6
<i>Longidorus</i> spp	15	15	1234±65	1
<i>Xiphinema</i> spp	15	15	1234±65	1

\*  $n/N \times 100$  (n = number of times individual nematodes occurred and N = Sample size (99)).

\*\*  $In/N \times 100$  (n = Individual nematode in all the samples and TN = Total Population of all the nematodes extracted in all the samples).

**Table 6.** Plant Parasitic Nematodes extracted from the roots of Yam in Oyo State.

Nematode Genera	Frequency of Occurrence	% Frequency Rating *	Nematode Population/10 g root	% Nematode Population **
<i>Meloidogyne</i> spp	64	65	8765±50	25
<i>Scutellonema</i> spp	59	60	6385±45	18.2
<i>Pratylenchus</i> spp	59	60	5836±45	16.3
<i>Aphelenchoides</i> spp	40	40	3283±35	9.3
<i>Trichodorus</i> spp	32	32	2355±35	6.4
<i>Radopholus</i> spp	30	30	2056±30	6
<i>Helicotylenchus</i> spp	25	25	1685±28	5
<i>Rotylenchus</i> spp	15	15	1668±22	5
<i>Longidorus</i> spp	10	10	1556±16	4.4
<i>Xiphinema</i> spp	10	10	1556±16	4.4

\* n/N x 100 (n = number of times individual nematodes occurred and N = Sample size( 99).

\*\* In/N x 100 (n = Individual nematode in all the samples and TN = Total Population of all the nematodes extracted in all the samples).

change and lack of stability in the ecosystem, making conditions favorable for certain species of plant parasitic nematodes while exerting immense selection pressure upon others (Wallace, 1971; Tiyaqi et al., 1987). The sustainability of these intensified yam based systems is threatened by build up of soil borne constraints, particularly plant parasitic nematodes.

Plant parasitic nematodes have been reported to constitute serious impediments to yam production in various parts of the world (Acosta and Ayala, 1975; Bridge, 1972; Thompson et al., 1973; Coates- Beckford and Braithwaite, 1977; Adesiyani and Odihirin, 1977; Caveness, 1982; Hahn et al., 1989; Weber et al., 1995). The findings of this investigation are similar to an earlier report by Unny and Jerath (1965) who identified eleven genera of plant parasitic nematodes associated with yam in the former Eastern Nigeria. In their study they identified *Scutello-nema* spp and *Meloidogyne* spp as the most important nematodes limiting yam production. Caveness (1965) found 28 plant parasitic nematodes associated with yams in West and Mid-West, East and North of Nigeria and listed four genera, *Scutellonema* spp., *Pratylenchus* spp., *Meloidogyne* spp and *Rotylenchulus* spp as being very important. Adesiyani and Odihirin (1977) identified six genera of plant parasitic nematodes in Mid-West State of Nigeria. The species identified were *Scutellonema* spp., *Pratylenchus* spp., *Meloidogyne* spp., *Helicotylenchus* spp., *Criconemoides* spp and *Xiphinema* spp. Acosta and Ayala (1975), Coates-Beckford and Braithwaite (1977) and Bridge and Page (1984) had made similar observations in Puerto-Rico, Trinidad and Papua New Guinea respectively.

Plant-parasitic nematodes damage is an important factor in tuber quality reduction and yield loss in yams both in the field and in storage. Yams are vulnerable to nematode damage as they reduce the yield and quality of the tubers as a result of root gallings, root lesions, dry

and soft rots depending on the type of plant parasitic nematodes present.

The presence of plant parasitic nematodes could constitute serious impediments to the growth and yield of yams in Edo, Ekiti and Oyo States.

Enlightenment Programme for the yam farmers should therefore be embarked upon by the State Agricultural Development Programmes in the zone to inform the farmers of the presence of plant parasitic nematodes in their farms and attendant implications.

## ACKNOWLEDGEMENT

We wish to thank the Director Professor E. A. Adebo-wale, Coordinator for RTEP Professor T. A. Akinlosotu, Extension Staff of Edo, Ekiti and Oyo State ADPs and Dr. A.A. Adeniji Project Manager, Root and Tuber Expansion Programme, Ijebu-Ife, Ogun State for supporting the project financially.

## REFERENCES

- Acosta N, Ayala A (1975). Pathogenecity of *Pratylenchus coffeae*, *Scutellonema bradys*, *Meloidogyne incognita* and *Rotylenchulus reniformis* on *Dioscorea rotundata*. J. Nematol. 7(1): 1-6.
- Adesiyani SO, Odihirin RA (1977). Plant -parasitic nematodes associated with yam tubers in Mid-West State, Nigeria. Nig. J. Plant Prot. 3: 171-179.
- Adegbite AA, Adesiyani, SO, Agbaje GO, Omoloye AA (2005). Host Suitability of Crops under Yam Intercrop to Root-knot Nematode (*Meloidogyne incognita* Race 2) in South-Western Nigeria. J. Agric. Rural Develop. Trop. Subtrop. 106 (2): 113-118.
- Agbaje GO, Adegbite AA, Akinlosotu TA (2003). Performance of new hybrid yam (*D. rotundata* Poir) varieties in the forest zone of Nigeria. Tropicultura 21(3): 149-152.
- Agbaje GO, Adegbite AA, Akinlosotu TA, Shoyinka SA (2002). Performance of new hybrid yam varieties (*D. rotundata* Poir) under different cropping systems. Afr. J. Root and Tuber Crops 5 (1): 8-11.
- Ayala A, Acosta N (1971). Observations on Yam (*D. alata*) nematodes. Nematropica 1(2): 39 – 40.
- Bell M (2004). Plant Parasitic Nematodes: Lucid key to 30 Genera of Plant Parasitic Nematodes. <http://www.lucidcentral.com/keys/nemato->

- todes.
- Bridge J (1972). Nematode problems with yams. *Dioscorea spp.* in Nigeria. Pest Articles and News summaries 18(1):89–91.
- Bridge J, Page SLJ (1984). Plant nematode pests of crops in Papua New Guinea. J. Plant Protect. Trop. 1:99-109.
- Caveness FE (1965). End of tour report on the Nematology Project. Ministry of Agriculture and Natural Resources (MANR), Western Region. Nigeria and United States Agency for International Development (USAID) Project. 620-11-110-050. p. 135.
- Caveness FE (1982). Plant parasitic nematodes and IITA mandate food crops: A position paper. IITA, Ibadan, Nigeria. p.15.
- Coates-Beckford, PL, Braithwaite, CWD, (1977) Comparison of various treatments for the control of *P. coffeae* in yam. Nematropica 7 (2): 20-26.
- Coursey DG (1967). Yams. Longmans. London p. 230.
- FAOSTAT data (2004). Food and Agriculture Organization of the United Nations, Rome.
- Green KR, Florini DA (1996). Pests and Pathogens of Yams in Storage. A meeting report. Afr. J. Root and Tuber Crops 1(2):38–42.
- Hahn SK, Isoba JCG, Ikotun T (1989). Resistance Breeding in Root and Tuber crops at the IITA, Ibadan, Nigeria. Crop Protect. 8:147 – 168.
- Nweke FI, Ugwu BO, Asadu CLA, Ay P (1991). Production costs in the yam-based cropping systems of southwestern Nigeria. Research Monograph No. 6, IITA, Ibadan, Nigeria. Resource and Crop Management Division. p.29.
- Ricka DA, Barker KR (1992). Nematode Assays and Advisory Services. In: Nematology in the Southern Region of the United States R. D. Riggs (Ed) South Cooperative Service Bulletin 276 Arkansas Agric. Exp. Fayetteville, Arkansas. pp. 8-20.
- Southey JF (1986). Laboratory methods for work with plant and soil nematodes. Reference Book Ministry of Agriculture Fisheries and Food No. 402. London, Her Majesty's Stationery Office. p. 202.
- Thompson AK, Been BO, Perkins C (1973). Nematodes in Stored Yams. Exp. Agric. 9(3): 281 – 286.
- Tiyagi SA, Bano M, Anver S (1987). Populations build up of phytonematodes associated with pearl millet. Int. Nematol. Network Newslett. 4 (4): 19-20.
- Unny KL, Jerath ML (1965). Parasitic nematodes on *Dioscorea spp.* in Eastern Nigeria. Plant Dis. Rep. 49: 875-876.
- Wallace HR (1971). The influence of density of nematode populations on plants Nematologica 17: 154 – 166.
- Weber G, Chindo PS, Elemo KA, Oikeh S (1995). Nematodes as production constraints in intensifying cereal-based cropping systems of the Northern Guinea Savannah. Resource and Crop Management Research Monograph No. 17. I.I.T.A. Nigeria. p. 36.