

## Review

# Utilization of fungi for biotreatment of raw wastewaters

COULIBALY Lacina<sup>1\*</sup>, GOURENE Germain<sup>1</sup>, AGATHOS N Spiros<sup>2</sup>

<sup>1</sup>Laboratoire d'Environnement et de Biologie Aquatique (LEBA), UFR-Sciences et Gestion de l'Environnement, Université d'Abobo-Adjamé, 02 BP 801 Abidjan 02, Côte d'Ivoire <sup>2</sup>Unit of Bioengineering, Catholic University of Louvain, Place Croix du Sud 2 Bte 19, 1348 Louvain-la-Neuve, Belgium

Accepted 21 April, 2019

Fungal biomasses are capable of treating metal-contaminated effluents with efficiencies several orders of magnitude superior to activated carbon (F-400) or the industrial resin Dowex-50. Additionally, fungal biomasses are susceptible to engineering improvements and regeneration of their capabilities. With regard to organic pollutants, excessive nutrients and dyes, fungi can remove them from wastewaters, leading to a decrease in their toxicities. However, the detoxification rates seem to be dependent on media and culture conditions. The posttreatment by anaerobic bioprocesses of effluents that have been pretreated with fungi can lead to higher biogas than the original effluents. In addition to the degradation of organic pollutants, fungi produce added-value products such as enzymes (LiP, MnP, Lacc, amylase, etc.) and single-cell protein (SCP). Most research on fungal capacities to purify polluted effluents has been performed on a laboratory scale, hence there is a need to extend such research to pilot scale and to apply it to industrial processes.

**Key words:** Wastewaters, effluents, fungi, biodegradation, biosorption, decolourisation, value-added treatment.

## INTRODUCTION

Fungi are recognized for their superior aptitudes to produce a large variety of extracellular proteins, organic acids and other metabolites, and for their capacities to adapt to severe environmental constraints (Lilly and Barnett, 1951; Cochrane, 1958). For example, *Aspergillus niger* is the prototypical fungus for the production of citric acid (Clark, 1962; Lal, 1980; Grewal and Kalra, 1995), homologous proteins (esp. enzymes) and heterologous proteins (Archer et al., 1994; Prasertsan et al., 1997; Radzio and Kuck, 1997; Wongwicharn et al., 1999; Xu et al., 2000). Moreover,

*Phanerochaete chrysosporium* is the model of white-rot fungi for the production of peroxidases (Bumpus et al., 1985; Rodriguez et al., 1999). Beyond the production of such relevant metabolites, fungi have been attracting a growing interest for the biotreatment (removal or destruction) of wastewater ingredients such as metals, inorganic nutrients and organic compounds (Akthar and Mohan, 1986; Field et al., 1993; Feijoo and Lema, 1995; Palma et al., 1999; Coulibaly, 2002).

The focus of this review therefore concerns the use of fungi to remove or degrade various wastewater constituents. Some instances of synthetic wastewaters are reported, but only the contributions of fungal biomass in the biological treatment of raw wastewater are discussed in some length.

Corresponding author. E-mail: [lacina91@hotmail.com](mailto:lacina91@hotmail.com). Phone: 00 (225) 07 49 71 53.

**Abbreviations;** SCP: single-cell protein, LiP: lignin peroxidase, MnP: manganese peroxidase, MIP: manganese independent peroxidase, Lacc: laccase, COD: chemical oxygen demand, HTL: heat treatment liquor, BOD: biochemical oxygen demand, OMW: olive mill wastewaters, WWTP: wastewater treatment plant.

## DOMESTIC SEWAGE

Domestic sewage contains carbon and nutrient sources that can be removed by fungal biomass. In an early

investigation, Thanh and Simard (1973) demonstrated the capacities of seventeen fungal biomasses to remove phosphates (84.1%), ammonia (73.3%), total nitrogen (68.1%) and chemical oxygen demand (COD) (39.3%). They obtained fungal growth on this effluent with an accumulation of biomass ( $451.2 \text{ mg l}^{-1}$ ) that contained protein ( $47\% \text{ g g}^{-1}$ ). There was variability in fungal capacities as to the removal of pollutants (see Table 1). In fact, *Trichothecium roseum* was the best in phosphate removal (97.5%), whilst *Epicoccum nigrum*, *Geotrichum candidum* and *Trichoderma sp.* were the best in the removal of ammonia (84%), total nitrogen (86.8%) and COD (72.3%), respectively. Concerning cell-protein production, *Paecilomyces carneus* had the highest ratio of protein to biomass (92.5%). However, this fungus did not grow very well on domestic sewage. In our laboratory, domestic wastewater pretreatment by a strain of *A. niger* has been investigated under transient conditions. This fungal biomass removed about 72% of COD and 65% of protein (Coulibaly, 2002). Despite the differences between the bioprocess investigated in these two studies, COD and protein removal rates are in the same order. The overall feasibility of domestic wastewater treatment under sewer -simulating conditions has been explored recently both experimentally and by simulation (Coulibaly, 2002; Coulibaly et al., 2002; Coulibaly and Agathos, 2003). The heat treatment liquor (HTL) of an activated sludge was decolourised by *Coriolus hirsutus* (Fujita et al., 2000). This fungal strain exhibited a strong ability to decolourise HTL (70%) with an accumulation of manganese independent peroxidase (MIP) and manganese peroxidase (MnP). Optimising the culture medium by adding nitrogen and carbon sources and improving the biomass quality resulted in increased colour removal capacity by *C. hirsutus* (Kumar et al., 1998; Miyata et al., 2000; Fujita et al., 2000). Although fungal applications have shown good capacities on sewage treatment, they are still underutilised in practice. This could be explained, in part, by a widespread *a priori* assumption that fungal strains do not perform as well as bacteria.

## AGROINDUSTRIAL EFFLUENTS

Industries of olive oil, tapioca starch, distillery (molasses), cotton bleaching, pulp and paper processing produce several billion litres of coloured, often toxic and harmful wastewaters over the world annually. Those effluents have strong concentrations of COD ( $10\text{-}200 \text{ g l}^{-1}$ ), phenol and its derivatives ( $0.5\text{-}8 \text{ g l}^{-1}$ ) and often contain proteins, cyanides, chlorinated lignin compounds and dyes (Borja et al., 1992, 1997; Nieto et al., 1992; Bengtsson and Triet, 1994; Garcia et al., 1997; Jimenez and Borja, 1997; Yesilada et al., 1998; Kahmark and Unwin, 1999). The large amount of lignin derivatives of these effluents is responsible of their dark-brown colour (Calvo et al.,

1995). The phenolic compounds of such wastewaters exert some bactericidal effects on wastewater treatment plant (WWTP) microorganisms (Borja et al., 1996; Fang and Chan, 1997; Vassilev et al., 1997; Sayadi et al., 2000). Fungal pretreatment (Table 2) of these effluents under aerobic conditions makes it possible to obtain phenol reduction (51-100%), good decolourisation (31-100%), biochemical oxygen demand (BOD) reduction up to 85.4%, and enzyme production (protease, Lacc (EC 1.10.3.2); LiP (EC 1.11.1.14), MnP (EC 1.11.1.13), amylase, etc.) (Vinciguerra et al., 1995; Yesilada et al., 1995; Garcia et al., 1997, 2000; D'Annibale et al., 1998; Setti et al., 1998; Gharsallah et al., 1999; Robles et al., 2000; Kissi et al., 2001). Amendment of olive mill wastewater (OMW) composition (addition of co-substrate, nutrients, salts) influences the removal of COD, phenols and colours (Yesilada et al., 1998). In fact, Garcia et al. (2000) noted that *G. candidum* removed COD but did not degrade phenols. However, by optimising OMW composition (COD:N:S = 100:5:2) for *G. candidum* growth, Assas et al. (2000) obtained a complete degradation of phenols and 70% decolourisation. Miranda et al. (1996) maximized colour removal from molasses wastewaters (up to 69%) with *A. niger*, after the amendment of the culture medium with co-substrate and mineral nutrients ( $\text{MgSO}_4$ ,  $\text{KH}_2\text{PO}_4$  and  $\text{NH}_4\text{NO}_3$ ). Some of the consequences of OMW pretreatment by fungi are the 23- to 30-fold higher increases in biogas production and the fertilizing effect on plants (*Trifolium repens*) compared to non-pretreated effluent (Borja et al., 1993, 1995 a,b,c; Jimenez and Borja, 1997; Vassilev et al., 1998).

The influence of co-substrate (see Table 3) upon paper and pulp industrial wastewater treatment, detoxification and decolourisation rates has also been observed with *Ceriporiopsis subvermisporea*, *P. chrysosporium*, *Trametes versicolor*, *Rhizopus oryzae* and *Rhizomucor pusillus* (Manzanares et al., 1995; van Driessel and Christov, 2002; Nagarathnamma and Bajpai, 1999; Nagarathnamma et al., 1999). The mechanisms of decolourisation of agroindustrial effluents by fungi are reported to include biosorption and/or biodegradation (Ohmomo, 1988; Sayadi and Ellouz, 1995; Soares and Duran, 1998; Christov et al., 1999; Nagarathnamma et al., 1999). Some "mycoreactors" such as rotating biological contactor (MYCOR), trickling filter reactor (MYCOPOR) and continuous column reactor have been developed to decolourise pulp and paper wastewaters (Eaton et al., 1982; Messner et al., 1990; Bajpai et al., 1993). These reactors were able to run over several weeks by maintaining their colour removal rates. Ligninolytic enzymes are also involved in the degradation of organic compounds, including dyes (see below), within these effluents (Chivukula et al., 1995). The enzymatic oxidation mechanism of those pollutants has been well discussed elsewhere and is not the aim of this contribution (Young and Yu, 1997; Mester and Tien,

**Table 1.** Examples of fungi used to treat domestic sewage, starch processing and metal bearing effluents. Optimal culture condition and the effect of fungal pretreatment are reported.

Effluents	Fungi	Treatment		References
		Reactor and medium handling	Parameters	
Domestic sewage	<i>Penicillium citricum</i> , <i>Steganosporium piriforme</i> , <i>Arthrinium arundis</i> , <i>Fusarium oxysporum</i> , <i>Cladosporium herbarum</i> , <i>Cladosporium cladosporioides</i> , <i>Scopulariopsis brevicaulis</i> , <i>Mucor hiemalis</i> , <i>Trichothecium roseum</i> , <i>Epicoccum nigrum</i> , <i>Helminthosporium sativum</i> , <i>Ulocladium atrum</i> , <i>Geotrichum candidum</i> , <i>Trichocladium asperum</i> , <i>Paecilomyces carneus</i> , <i>Trichoderma sp.</i> , <i>Chrysosporium pannorum</i>	Shake-flask	COD (72.3%); Phosphates (97.5%); N-total (86.8%); Dry matter (684 mg l <sup>-1</sup> ); Protein content (205 mg l <sup>-1</sup> )	Thanh and Simard (1973)
	<i>Aspergillus niger</i>	Stirred tanks reactor in series	COD (72%); N-total (65.4%)	Coulibaly (2002)
	<i>Corioliolus hirsutus</i>	Continuous immobilized bioreactor; addition (nutrient (NH <sub>4</sub> (100 mg l <sup>-1</sup> ), NO <sub>3</sub> (100 mg l <sup>-1</sup> ); MnSO <sub>4</sub> ); co-substrate (glucose, 0.5%)	Decolorization (80%, 2 d); MnP (60 U l <sup>-1</sup> ); MIP (40 U l <sup>-1</sup> )	Miyata et al. (2000)
Starch process effluent	<i>A. oryzae</i> ; <i>Rhizopus arrhizus</i> ; <i>Trichoderma viride</i> ; <i>T. reesei</i> ; <i>G. candidum</i> ; <i>A. terreus</i> ; <i>R. oligosporus</i>	Shake-flask, air lift bioreactor (45 l); addition of nutriment (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> ; Urea; NH <sub>4</sub> NO <sub>3</sub> ; NaNO <sub>3</sub> ; K <sub>2</sub> HPO <sub>4</sub> ; KH <sub>2</sub> PO <sub>4</sub> )	TOC (44-88%); SS (95%); starch hydrolysis (53-100%); biomass (2-5.6 g l <sup>-1</sup> ); protein (48.8% of biomass weight); COD (97.8%); glucoamylase (3.94 U ml <sup>-1</sup> )	Jin et al. (1999abc; 2001)
	<i>A. niger</i> ; <i>A. oryzae</i>	Shake-flask	COD (90%); biomass and amylase production	Fujita et al. (1993); Murado et al. (1993)
Metal bearing effluent	<i>A. niger</i> , <i>P. simplicissimum</i> , <i>Geotrichum sp.</i> , <i>Fusarium verticillioides</i> , <i>Rhizoctonia solani</i> , <i>Aquathanatephorus pendulus</i> ;	Shake-flask, presence of co-ions, biomass (produced)	<i>A. niger</i> (Cu (91%); Zn (70%))	Price et al. (2001); Gomes et al. (1998, 1999); Gomes and Linardi (1996); Karavaiko et al. (1996)
	<i>A. niger</i> , <i>A. flavus</i> , <i>A. fumigatus</i> ; <i>R. Arrhizus</i> ; <i>A. terrus</i>	Shake-flask; presence of co-ions, biomass (industrial waste, produced)	Metal removal (82-100%)	Balakrishnan et al. (1994); Niyogi et al.(1998)
	<i>Mucor meihii</i>	Shake-flask; biomass (industrial waste), dilution (1-20)	Sorption (0.7-1.15 mmol g <sup>-1</sup> )	Tobin and Roux (1998)
	<i>A. niger</i>	Shake-flask; presence of co-ions, biomass (produced)	Metal removal (75%)	Akthar and Mohan (1995)

**Table 2.** Examples of fungi used to treat distillery wastewaters. Optimal culture condition and the effect of fungal pretreatment are reported.

Effluents	Fungi	Treatment		References
		Reactor and medium handling	Parameters	
Distillery Wastewaters	<i>A. awamori</i> var. <i>kawachi</i>	Shake-flask; Jar-fermentor (30 l)	Specific resistance of culture broth (97.5% decrease); BOD (56%); TOC (72%); Phosphates (80%) protein in mycelium (40%) Increased TOC removal of anaerobic pretreated effluent	Kida et al. (1995)
	<i>A. niger</i> ; <i>A. awamori</i>	Shake-flask; continuous bubble reactor; co-substrate (sucrose, fructose, glucose); MgSO <sub>4</sub> (1 g l <sup>-1</sup> ); KH <sub>2</sub> PO <sub>4</sub> (0.5 g l <sup>-1</sup> ); NH <sub>4</sub> NO <sub>3</sub> (1.8 g l <sup>-1</sup> ); peptone (5%); rice (3%)	OMW (decolorization (69%, 3-4 d); COD (78%)) Thin stillage (protease (200 U ml <sup>-1</sup> ); biomass (30 g l <sup>-1</sup> ))	Miranda et al. (1996); Yang and Lin (1998)
	<i>P. chrysosporium</i> ; <i>G. candidum</i> ; <i>C. versicolor</i> ; <i>Mycelia sterilia</i>	Dilution (50%)	Decolorization (53%, 10 d); growth rates inhibition below 50% of dilution; decoloration of melanoidins (80%) by <i>P. chrysosporium</i> JAG-40	FitzGibbon et al. (1998)
	<i>G. candidum</i>	Shake-flask; Jar-fermentor (7 l); Polyurethane-foam; immobilization; co-substrate (glucose, 0.5-1%)	Decolorization (80%), peroxidase accumulation	Kim and Shoda (1999)
	<i>Trametes versicolor</i>	Shake-flask; inoculum size, sucrose addition (0.3%); KH <sub>2</sub> PO <sub>4</sub> (0.5 g l <sup>-1</sup> );	Decolorization (82%); COD (77%); NH <sub>4</sub> <sup>+</sup> (36%)	Benito et al. (1997)
	<i>C. hirsutus</i>	Shake-flask; continuous immobilized polyurethane-foam reactor; MnSO <sub>4</sub> ; co-substrate (glucose, ethanol)	Decolorization (76%, 2 d); TOC (45%);	Miyata et al. (2000)
	<i>Flavodon flavus</i> ; <i>P. decumbens</i>	Shake-flask; aeration; co-substrates (sucrose, glucose, mannose, mannitol, xylose, arabinose, fructose, glycerol) tested at 10%	Decolorization (80%); MnP(400 U l <sup>-1</sup> ); Lacc (550 nkat l <sup>-1</sup> ); increase of anaerobic digestion rate of aerobic pretreated effluent by <i>P. decumbens</i>	Raghukumar and Rivonkar (2001); Jimenez and Borja (1997)
	<i>Ceriporopsis subvermispora</i>	Shake-flask; co-substrat (glucose, 0.1%; sucrose; lactose; microcrystalline cellulose; carboxymethyl cellulose; xylose; starch; athyl alcohol; bagasse pith; cheese whey; prehydrolysate liquor; molasses)	Color (90%); COD (45%); lignin (62%); AOX (32%); EOX (36%)	Nagarathnamma et al. (1999)

**Table 3.** Examples of fungi used to treat wood processing wastewaters. Optimal culture condition and the effect of fungal pretreatment are reported.

Effluents	Fungi	Treatment		References
		Reactor and medium handling	Parameters	
Wood processing wastewaters	<i>Sporotichum pulverulentum</i> ( <i>P. chrysosporium</i> )	Batch reactor (25 m <sup>3</sup> ); continuous laboratory fermentor (10 l)	Biomass (5.7 g l <sup>-1</sup> ); Protein (42%); Protein productivity (132 mg l <sup>-1</sup> ); cellulase (0.2 U ml <sup>-1</sup> ); Suspended particles (88%); BOD 73%; COD 52%); feedstuff (rat, pigs and sheep)	Ek and Eriksson (1980); Thomke and Rundgreen (1980)
	<i>A. foetidus</i>	Shake-flask; dilution (10%)	Decolorization (90%, 2 d)	Sumathi and Phatak (1999)
	<i>C. versicolor</i> ; <i>P. chrysosporium</i> ; <i>Pleurotus ostreatus</i> ; <i>polyporus versicolor</i>	Shake-flask; immobilization in beads of Ca-alginate gel; dilution (16.7%); co-substrates (sucrose 0.5%, xylose, glucose, glycerol, ethanol)	Decolorization of suspended biomass (60%, 6 d); decolorization of immobilized biomass (80%, 3 d)	Livernoche et al (1980); Marwaha et al. (1998)
	<i>P. chrysosporium</i> ; <i>Phanerochaete flavid-alba</i>	Shake-flask; co-substrate (glucose); Mn (0.3 mg l <sup>-1</sup> ); culture age	Decolorization (88%), LiP (450 nmol min <sup>-1</sup> ml <sup>-1</sup> ); MnP (800 nmol min <sup>-1</sup> ml <sup>-1</sup> )	Perez et al. (1997)
	<i>P. chrysosporium</i> ; <i>Ganoderma australe</i> ; <i>Coriopsis gallica</i> ; <i>Paecilomyces variotii</i>	Shake-flask	<i>P. chrysosporium</i> (decolorization, 50%, 7 d; lignin pyrolysis compounds, 57% reduction); <i>G. australe</i> (decolorization, 50%, 7 d; lignin derivated compounds 48% increased); <i>C. gallica</i> (decolorization, 48%, 7 d; lignin-derivated compounds 77% reduction); <i>P. variotii</i> (decolorization, 85%, 7 d; lignin-derivated compounds 78% reduction)	Calvo et al. (1995 a b)
	<i>T. versicolor</i>	Shake-flask; continuous feeding bioreactor; culture age; dilution 30%); SO <sub>4</sub> Mn (23 mg l <sup>-1</sup> ); co-substrate (glucose, 0.3%; sucrose; starch; ethanol, carboxymethyl-cellulose; pulp and bagasse pith)	Decolorization (90 %, 9 d) Lacc (700 U l <sup>-1</sup> , 10 d); MnP (25 U l <sup>-1</sup> , 7 d); phenols (90%); COD (69%)	Manzanares et al. (1995); Mehna et al. (1995)
	<i>Sordaria fimicola</i> ; <i>Halosarpeia ratnagiriensis</i>	Shake-flask; pH (4.5; 8.2)	<i>S. fimicola</i> (decolorization, 55%, Lacc (100 nkatal ml <sup>-1</sup> )); <i>H. ratnagiriensis</i> (decolorization, 85%; Lacc (100 nkatal ml <sup>-1</sup> ))	Raghukumar et al. (1996)
	<i>Ceriporopsis subvermispota</i> ; <i>R. oryzae</i>	Shake-flask; co-substrat (glucose, 0.1%; sucrose; lactose; microcrystalline cellulose; carboxymethyl cellulose; xylose; starch; athyl alcool; bagasse pith; cheese whey; prehydrolysate liquor; molasses)	Color (90%); COD (45%); lignin (62%); AOX (32%); EOX (36%)	Nagarathnamma et al. (1999); Nagarathnamma and Bajpai (1999)

2000). Beneficial effects of the fungal pretreatment of pulp mill effluent upon its subsequent anaerobic digestion have been reported (Feijoo et al., 1995). Anaerobic digestion of Kraft pulp mill effluent pretreated by *P. chrysosporium* gave increased degradation of high molecular weight compounds (79%) according to these authors. Also, an important decolourisation (79%) was also observed, that was correlated with MnP accumulation.

With regard to other agroindustrial wastewaters that are relatively non toxic (e.g. dilute lignocellulosics, starch, rice and mussels processing, sauce production, etc.) (see Table 1), fungal growth on them has been reported to produce single-cell protein (SPC), enzymes, chitosan, amyolytic preparations and a good reduction of COD (up to 97.8%) (Morimura et al., 1992, 1994 a,b; Murado et al., 1993; Kida et al., 1995; Yang and Lin, 1998; Yokoi et al., 1998; Jin et al., 1998, 1999, 2001).

## DYED EFFLUENTS

The effluents of pharmaceutical industries, dyeing, printing, photographs, textile and cosmetics contain dyes (McMullan et al., 2001). For example, over  $7 \times 10^7$  tons dyes are produced annually worldwide, of which about 10% are lost in industrial effluent (Vaidya and Datye, 1982). Wastewaters from textile industries are a complex mixture of many polluting substances such as organochlorine-based pesticides, heavy metals, pigments and dyes. Their compositions have been discussed in detail by O'Neill et al. (1999). The majority of these dyes are slowly removed by the WWTP, because of their toxicities to indigenous microorganisms. Dye removal from wastewaters by established WWTP processes are expensive and need careful application (Vandevivere et al., 1998; Robinson et al., 2001). Furthermore, following anaerobic digestion, nitrogen-containing dyes are transformed into aromatic amines that are more toxic and mutagenic than the parent molecules (Shaul et al., 1985; Chung and Stevens, 1993; Ganesh et al., 1994). To overcome these difficulties, fungi are being investigated for their potential to decolourise effluents. Among them, the most widely studied are the white-rot fungi *P. chrysosporium* (a model, primarily laboratory organism) and *T. versicolor* (a promising organism for industrial applications).

Nowadays other fungi have also shown some capacities to remove dyes from industrial effluents. Dyes are removed by fungi by biosorption (Contato and Corso, 1996; Tatarko and Bumpus, 1998; Payman et al., 1998; Zheng et al., 1999; Fu and Viraraghavan, 2000), biodegradation (Nigam et al., 1995; Conneely et al., 1999) and enzymatic mineralisation (LiP, MnP, manganese independent peroxidase (MIP), Lacc) (Young and Yu, 1997; Ferreira et al., 2000; Ollikka et al., 1998; Podgornik et al., 1999; Wong and Yu, 1999; Zheng et al.,

1999; Pointing and Vrijmoed, 2000; Wesenberg et al., 2003). However, one or more of these mechanisms could be involved in colour removal, depending on the fungus used. Other fungal biomasses applied to the decolourisation of raw textile effluents include *Botrytis cinerea*, *Endothiella aggregata*, *Geotrichum fici*, *R. oryzae*, *Tremella fuciformis*, *Xeromyces bisporus*, *Hirschioporus larincinus*, *Inonotus hispidus*, *Phebia tremellosa* and *C. versicolor* (Banat et al., 1996; Kirby, 2000; Polman and Breckenridge, 1996). It is reported that raw effluents can only partially be decolourised upon fungal treatment (maximum of 49-80% but often much less). For example, a complex mixture of real textile effluents containing many reactive dyes could be decolourised upon partial dilution by using the agaric white-rot fungus *Clitocybula duseinii* (Wesenberg et al., 2002). The weak decolourisation of these effluents by complete cultures could be explained by the influences of temperature, pH, salts, inhibitory molecules (sulphur compounds, surfactants, heavy metals, bleaching chemicals), carbon and nutrients within these solutions (Chao and Lee, 1994; Jacob et al., 1998; Swamy and Ramsay, 1999; Mester and Tien, 2000). Concerning enzymatic (Lacc, LiP, MnP) degradations, these reactions are quite complicated, involving numerous low molecular weight cofactors that serve as redox mediators (Reyes et al., 1999; Wesenberg et al., 2003). These cofactors, in addition to the enzymes themselves, influence fungal colour removal rates.

## METAL CONTAINING EFFLUENTS

Metallurgical industries, mining, surfaces cleaning, waste incinerators produce large wastewater polluted by metals. Dissolved metals escaping into the environment pose a serious health hazard. Because they accumulate in living tissues throughout the food chain, which has human at its top. There is a need to remove heavy metals before they enter the complex ecosystem. Physicochemical treatments evolved in very diluted water-containing metals (precipitation, electrochemical, flocculation, coagulation, ion exchange) are expensive. Utilization of biomasses in general (Volesky, 1994; Veglio and Beolchini, 1997; Kratochvil and Volesky, 1998; McKay et al., 1999; Gupta et al., 2000) and particularly that of fungi are considered to be best alternatives for those waters purification (Kapoor and Viraraghavan, 1995; Modak and Natarajan, 1995; Sag et al., 1998; Volesky and Holan, 1995; Atkinson et al., 1998; Kratochvil and Volesky, 1998; Mogollon et al., 1998; Savvaidis, 1998; Tobin and Roux, 1998). Indeed, the purification of the water-containing metals by fungal biomass is cheaper and it presents the following advantages: (i) production of residual small volume; (ii) possibility of valorisation of fungal waste biomasses from industrial fermentations; (iii) fast removal and (iv) easy installation of the process.

Fungal biomasses walls are composed of macromolecules (chitin, chitosan, glucan, lipid, phospholipides), which contain carboxyl groups (R-COOH), amino groups (R<sub>2</sub>NH, R-NH<sub>2</sub>), phosphates, lipids, melanin, sulphates (R-OSO<sub>3</sub><sup>-</sup>) and hydroxides (OH<sup>-</sup>) (Caesartonthat et al., 1995; Kapoor and Viraraghavan, 1998 a,b; Fogarty and Tobin, 1996; Kapoor et al., 1999). Those functional groups are metals sorption sites (Tsezos and Volesky, 1982; Mullen et al., 1992; Guibal et al., 1995; Gardea Torresdey et al., 1996; Kapoor and Viraraghavan, 1997; Matheickal and Yu, 1997; Zhang et al., 1998; Sarret et al., 1999; McHale and McHale, 1994; Mashitah et al., 1999; Tereshina et al., 1999; Zhou, 1999). Fungi remove metals essentially by adsorption, chemisorptions (ion exchange), complexation,

coordination, chelation, physical adsorption and microprecipitation (Guibal et al., 1995; Huang and Huang, 1996; Kapoor and Viraraghavan, 1997; Sarret et al., 1998). There are also possible oxydo-reduction taking place in the biosorbent. When metals are removed by ionic exchange, they generally replace K<sup>+</sup>, Mg<sup>2+</sup>, Ca<sup>2+</sup> and H<sup>+</sup> contained in biomasses (Akthar et al., 1995; Zhou, 1999; Gomes et al., 1999; Mashitah et al., 1999). Table 1 gives a synthesis of some works on metals removal from wastewaters by some fungi. Biomasses used to remove metals from wastewaters are generally produced against few residual biomasses from fermentation (Fourest et al., 1994; Meyer and Wallis, 1997). Metals sequestrations by fungi are influenced by the mineral and organic compositions content of the medium in which biomasses are produced. Biomasses granulometries and physiological states (living or dead), co-ions, metals concentrations and physical parameters (temperature, pH, ionic force, presence of others metals) influence also metals removal from polluted waters (Volesky, 1994; Akthar et al., 1995; Gomes and Linardi, 1996; Modak et al., 1996; Gardea Torresdey et al., 1997; Yu and Kaewsan, 1999; Zhou, 1999). Metals by fungi from various raw effluents (gold mining effluent, tanning effluent, swine water, polluted lake waters) are sometimes completely removed (see Table 1). However, these outputs depend on the metal and fungus involved.

To increase fungal biomasses removal capacities, some of them undergo physicochemical treatments (soda or acidic treatments, insertion of functional groupings, heat treatment) (Akthar et al., 1995; Akthar and Mohan, 1995; Kapoor et al., 1999; Kramer and Meisch, 1999; Yin et al., 1999; Yan and Viraraghavan, 2000). Moreover, *A. niger* biomass treatment by soda, makes it possible to adsorb 2.5 to 1000 mg Ag l<sup>-1</sup> of Ag<sup>+</sup> in polluted water (Akthar et al., 1995). In the same way, Kapoor et al. (1999) obtained with a soda treated biomass of *A. niger*, the removal rates of Cd<sup>2+</sup>, Cu<sup>2+</sup> and Pb<sup>2+</sup> superiors to that of activated carbon (F-400). Akthar and Mohan (1995) used the same type of biomass like precedent authors, and they obtained the removal rates of Zn<sup>2+</sup> and Cd<sup>2+</sup> superiors to that of Dowex-50 resin. An insertion of

carboxyl and amino groups in *A. niger* biomass walls, makes it possible to obtain an adsorption rate ranging in the order of 172-1064 mmol (kg biomass)<sup>-1</sup> for Cd<sup>2+</sup>, Co<sup>2+</sup>, Ni<sup>2+</sup> and Zn<sup>2+</sup> (Kramer and Meisch, 1999). A simple detergent and alkaline solutions treatment of *M. rouxii* biomass was sufficient to obtain an increase in its adsorption capacity (Yan and Viraraghavan, 2000). Fungal biomasses that have sequestered metals can be regenerated following their washing with HNO<sub>3</sub> (0.05 N) and/or with Ca<sup>2+</sup>, Mg<sup>2+</sup> and K<sup>+</sup> (0.1 M) (Akthar et al., 1995, 1996; Kapoor et al., 1999).

## DISCUSSION

Essential works on fungal utilization for raw wastewaters biopurification have been laboratory tests. This situation can be explained by the fact that fungal utilization in environmental biotechnology is still under investigation to assess information's on process implementation. To gain confidence with the results, these investigations are performed on synthetics wastewaters. The good results obtained in laboratory tests depend on growth medium optimisation (addition of co-substrate, nutrients, mediators molecules, physical parameters optimisation) and a good handling of biomasses. However, these works amongst other things prove some advantages when a mycoreactor is introduced in effluents treatment lines. In fact, there are some reductions of bactericidal effects and an increase in biogas production. Consequently work on pilot and the development of treatment plants are to be encouraged.

The degradation and the mineralisation of some recalcitrant dyes and organochlorinated compounds are effective by certain white rot fungi. However fungal aptitudes for raw wastewaters remain dependent on salts concentration, culture conditions and especially on the amendment of carbon and nutriment sources. Among the co-susubstrates tested for effluent pretreatment by fungi, glucose and sucrose were the best, when they were used at 5 to 10 g l<sup>-1</sup>. To minimise the mycoreactor integration cost in the treatment line, the co-substrate could be provided by feeding the reactors with amylaceous effluents or others wastewaters rich in sucrose or glucose as these carbon sources proved to be the best co-substrate. About the growth medium impact on fungal capacities to decolourise HTL, one could use *C. hirsutus* in post processing of an activated wastewater, because of its sensitivity to organic nitrogen. The oxidoreductases activities could be more significant, thanks to the use of substrate that could ensure the role of mediators' molecules and guarantee the generation of H<sub>2</sub>O<sub>2</sub> in the reaction medium. Salts constraint could be overcome while proceeding to the desalinisation of the effluents before their treatment with fungi.

As regards metal removal, a standardization of adsorption rates unit and the rigorous description of

biomass morphology (pellet diameter) will allow a better comparison of fungal capacities and a guide for the best choice of fungi. Biomass grinding to produce small particles and their engineering to increase their capacities to remove specific metals are promising way for fungal biosorption. Utilization of residual biomasses from fermentation is still minimal nowadays (Knapp and Newby, 1999); one needs to encourage such practice, because this constitutes a way of making use of them.

## CONCLUSION

This review highlighted the capacities of certain fungi to pretreat raw wastewaters. However, essential works on this subject are still laboratories tests and they are of less industrial scale application. The white rot fungi are suitable for the degradation of a large variety of pollutants and to produce at the same time metabolites of great added values (proteins, enzymes). However, an optimisation of the culture media in carbon sources or nutrients and mediators molecules is very important to obtain a good output of pollutants degradations. With regard to other fungi, those also contribute to effluents purification with proteins and enzymes productions (for example, *A. awamory* and *A. niger*). Some residual biomasses from fungal fermentation, have been used to remove metals and dyes from effluents. Ultimately, the fungal biomasses present many assets for biopurification of wastewaters.

## ACKNOWLEDGEMENTS

This research was supported by a fellowship of The Ministry of Higher Education, Scientific Research and Technological Innovation of Côte d'Ivoire. We thank all of our colleagues from LEBA and Unit of Bioengineering for their useful discussions.

## REFERENCES

- Akthar MN, Mohan PM (1995). Bioremediation of toxic metal ions from polluted lake waters and industrial effluents by fungal biosorbent. *Curr. Sci.* 69: 1028-1030.
- Akthar MN, Sastry KS, Mohan PM (1995). Biosorption of silver ions by processed *Aspergillus niger* biomass. *Biotechnol. Lett.* 17: 551-556.
- Akthar MN, Sastry KS, Mohan PM (1996). Mechanism of metal ion biosorption by fungal biomass. *Biometals.* 9: 21-28.
- Archer DB, Jeenes DJ, Mackenzie DA (1994). Heterologous protein production from filamentous fungi. *Antonie van Leeuwenhoek.* 65: 245-250.
- Assas N, Marouani L, Hamdi M (2000). Scale down and optimization of olive mill wastewater decolorization by *Geotrichum candidum*. *Bioprocess Eng.* 22: 503-507.
- Atkinson BW, Bux F, Kasan HC (1998). Consideration for application of biosorption technology to remediate metal-contaminated industrial effluents. *Water S.A.* 24: 129-135.
- Balakrishnan M, Modak JM, Natarajan KA, Naik JSG (1994). Biological uptake of precious and base metals from gold process cyanide effluents. *Min. Metall. Process.* 11: 197-202.
- Banat IM, Nigam P, Singh D, Marchant R (1996). Microbial decolorization of textile-dye-containing effluents: a review. *Biores. Technol.* 58: 217-227.
- Bengtsson BE, Triet T (1994). Tapioca starch wastewater toxicity characterized by Microtox and Duckweed tests. *Ambio.* 23: 473-477.
- Bhainsa KC, D'Souza SF (1999). Biosorption of uranium (VI) by *Aspergillus fumigatus*. *Biotechnol. Technique* 13: 695-699.
- Borja R, Martin A, Maestro R, Alba J, Fiestas JA (1992). Enhancement of the anaerobic digestion of olive mill wastewater by the removal of phenolic inhibitors. *Process Biochem.* 27: 231-237.
- Borja R, Garrido SE, Martinez L, RamosCormenzana A, Matin A (1993). Kinetic study of anaerobic digestion of olive mill wastewater previously fermented with *Aspergillus niger*. *Process Biochem.* 28: 397-404.
- Borja R, Alba J, Garrido SE, Martinez L, Garcia MP, Monteoliva M, RamosCormenzana A (1995a). Effect of aerobic pretreatment with *Aspergillus terreus* on the anaerobic digestion of olive mill wastewater. *Biotechnol. Appl. Biochem.* 22: 233-246.
- Borja R, Alba J, Garrido SE, Martinez L, Garcia MP, Incerti C, RamosCormenzana A (1995b). Comparative study of anaerobic digestion of olive mill wastewater (OMW) and OMW previously fermented with *Aspergillus terreus*. *Bioprocess Eng.* 13: 317-322.
- Borja R, Martin A, Alonso V, Garcia I, Banks CJ (1995c). Influence of different aerobic pretreatments on the kinetics of anaerobic digestion of olive mill wastewater. *J. Chem. Technol. Biotechnol.* 29: 489-495.
- Borja R, Banks CJ, Maestro-Duran R, Alba J (1996). The effects of the most important phenolic constituents of olive mill wastewater on batch anaerobic methanogenesis. *Environ. Technol.* 17: 167-174.
- Borja R, Alba J, Banks CJ (1997). Impact of the main phenolic compounds of olive mill wastewater (OMW) on the kinetics of acetoclastic methanogenesis. *Process Biochem.* 32: 121-133.
- Bumpus JA, Tien M, Wright D, Aust SD (1985). Oxidation of persistent environmental pollutants by white rot fungus. *Science.* 228: 1434-1436.
- Caesartonthat TC, Kloeke FV, Geesey GG, Henson JM (1995). Melanin production by filamentous soil fungus in response to copper and localization of copper sulfide by sulfide-silver staining. *Appl. Environ. Microbiol.* 61: 1968-1975.
- Calvo AM, Terron MC, Fidalgo ML, Pelayo JM, Galletti GC, Gonzalez AE (1995). Pyrolysis-gas chromatography-mass spectrometry characterization of wheat straw alkaline-cooking effluents after biological treatment with the fungi *Phanerochaete chrysosporium* and *Ganoderma australe*. *Analytica Acta* 309: 145-152.
- Chao WL, Lee SL (1994). Decoloration of azo dyes by 3 white rot fungi. Influence of carbon source. *World J. Microbiol. Biotechnol.* 10: 556-559.
- Chivukula M, Spadaro JT, Renganathan V (1995). Lignin peroxidase catalyzed oxidation of sulfonated azo dyes generates novel sulfophenyl hydroperoxides. *Biochem.* 34: 7765-7772.
- Christov LP, Van Driessel B, du Plessis CA (1999). Fungal biomass from *Rhizomucor pulvisillus* as adsorbent of chromophores from a bleach plant effluent. *Process Biochem.* 35: 91-95.
- Chung KT, Stevens SE Jr (1993). Decolorization of azo dyes by environmental microorganisms and helminthes. *Environ. Toxicol. Chem.* 12: 2121-2132.
- Clark DS (1962). Submerged citric acid fermentation of ferrocyanide-treated molasses: Morphology of pellets of *Aspergillus niger*. *Can. J. Microbiol.* 8: 133-136.
- Cochrane VW (1958) *Physiology of the fungi*. John Wiley and Sons Inc., New York, New York
- Conneely A, Smyth WF, McMullan G (1999). Metabolism of the phthalocyanine textile dye remazol turquoise blue by *Phanerochaete chrysosporium*. *FEMS Microbiol. Lett.* 179: 333-337.
- Contato M, Corso CR (1996). Studies of adsorptive interaction between *Aspergillus niger* and the reactive azo dye Procion Blue MX-G. *Eclat. Quim.* 21: 97-102.
- Coulibaly L (2002). Bioconversion de macromolécules dans un réacteur simulant un écoulement piston en régime transitoire. Cas de la bioremédiation d'eaux usées synthétique par *Aspergillus niger*. Thèse de doctorat, Université Catholique de Louvain, Unité de génie biologique. [www.gebi.ucl.ac.be](http://www.gebi.ucl.ac.be)
- Coulibaly L, Naveau H, Agathos SN (2002). A tanks-in-series bioreactor



- to simulate macromolecule-laden wastewater pretreatment under sewer conditions by *Aspergillus niger*. *Wat. Res.* 36: 3941-3948.
- Coulibaly L, Agathos SN (2003) Transformation kinetics of mixed polymeric substrates under transitory conditions by *Aspergillus niger*. *Afr. J. Biotechnol.* 2: 438-443.
- D'Annibale A, Crestini C, Vinciguerra V, Sermanni GG (1998). The biodegradation of recalcitrant effluents from an olive mill by a white-rot fungus. *J. Biotechnol.* 61: 209-218.
- Eaton DC, Chang HM, Joyce TW, Jeffries TW, Kirk TK (1982). Method obtains fungal reduction of the solar kraft bleach plant effluents. *Tappi.* 65: 89-92.
- Ek M, Eriksson KE (1980). Utilization of the white-rot fungus *Sporotrichum pulverulentum* for water purification and protein production on mixed lignocellulosic wastewaters. *Biotechnol. Bioeng.* 22: 2273-2284.
- Fang HHP., Chan OC (1997). Toxicity of phenol towards anaerobic biogranules. *Wat. Res.* 31: 2229-2242.
- Feijoo G, Lema JM (1995). Treatment of forest industry effluents with toxic and recalcitrant compounds by the white rot fungi. *Afinidad.* 52: 171-180
- Feijoo G, Vidal G, Moreira MT, Mendez R, Lema JM (1995). Degradation of high molecular weight compounds of kraft pulp mill effluents by a combined treatment with fungi and bacteria. *Biotechnol. Lett.* 17: 1261-1266.
- Ferreira VS, Magalhaes DB, Kling SH, da Silva JG, Bon EPS (2000). N-demethylation of methylene blue by lignin peroxidase from *Phanerochaete chrysosporium*: Stoichiometric relation for H<sub>2</sub>O<sub>2</sub> consumption. *Appl. Biochem. Biotechnol.* 84-86: 255-265.
- Field JA, de Jong E, Feijoo Costa G, de Bont JAM (1993) Screening for ligninolytic fungi applicable to the degradation of xenobiotics. *TiBtech.* 14: 44-49.
- FitzGibbon F, Singh D, McMullan G, Marchant R (1998). The effect of phenolic acids and molasses spent wash concentration on distillery wastewater remediation by fungi. *Process Biochem.* 33: 799-803.
- Fogarty RV, Tobin JM (1996). Fungal melanins and their interactions with metals. *Enzyme Microb. Technol.* 19: 311-317.
- Fourest E, Canal C, Roux JC (1994). Improvement of heavy metal biosorption by dead biomass (*Rhizopus arrhizus*, *Mucor miehei* and *Penicillium chrysogenum*) pH control and cationic activation. *FEMS Microbiol. Rev.* 14: 325-332
- Fu YZ, Viraraghavan T (2000). Removal of a dye from aqueous solution by the fungus *Aspergillus niger*. *Water Qual. Res. J. Can.* 35: 95-111.
- Fujita M, Iwahori K, Yamakawa K (1993). Pellet formation of fungi and its application to starch wastewater treatment. *Wat. Sci. Technol.* 28: 267-274.
- Fujita M, Era A, Ike M, Soda S, Miyata N, Hirao T (2000). Decolorization of heat-treatment liquor of waste sludge by a bioreactor using polyurethane foam-immobilized white rot fungi equipped with an ultramembrane filtration unit. *J. Biosc. Bioeng.* 90: 387-394.
- Ganesh R, Boardman GD, Michelsen D (1994). Fate of azo dyes in sludges. *Wat. Res.* 28: 1051-1057.
- Garcia IG, Venceslada JLB, Pena PRJ, Gomez ER (1997). Biodegradation of phenol compounds in vinasse using *Aspergillus terreus* and *Geotrichum candidum*. *Wat. Res.* 31: 2005-2011.
- Garcia IG, Pena PRJ, Venceslada JLB, Martin AM, Santos MAM, Gomez ER (2000). Removal of phenol compounds from olive mill wastewater using *Phanerochaete chrysosporium*, *Aspergillus niger*, *Aspergillus terreus* and *Geotrichum candidum*. *Process Biochem.* 35: 751-758.
- Gardea Torresdey JL, Cano Aguilera I, Webb R, Gutierrez Corona F (1996). Copper adsorption by inactivated cells of *Mucor rouxii*. *J. Hazard. Mater.* 48: 171-180.
- Gardea Torresdey JL, Cano Aguilera I, Webb R, Gutierrez Corona F (1997). Enhanced copper adsorption and morphological alteration of cells of copper-stressed *Mucor rouxii*. *Environ. Toxicol. Chem.* 16: 435-441.
- Gharsallah N, Labat M, Aloui F, Sayadi S (1999). The effect of *Phanerochaete chrysosporium* pretreatment of olive mill waste waters on anaerobic digestion. *Resources Conserv. Recycl.* 27: 187-192.
- Gomes NCM, Linardi VR (1996). Removal of gold, silver and copper by living and non living fungi from leach liquor obtained from the gold mining industry. *Rev. Microbiol.* 27: 218-222.
- Gomes NCM, Camargos ERS, Dias JCT, Linardi VR (1998). Gold and silver accumulation by *Aspergillus niger* from cyanide-containing solution obtained from the gold mining industry. *World J. Microbiol. Biotechnol.* 14: 149-149.
- Gomes NCM, Figueira MM, Camargos ERS, Mendonca-Hagler LCS, Dias JCT, Linardi VR (1999). Cyano-metal complexes uptake by *Aspergillus niger*. *Biotechnol. Lett.* 21: 487-490.
- Grewal HS, Kalra KL (1995). Fungal production of citric acid. *Biotechnol. Adv.* 13: 209-234.
- Guibal E, Roulph C, Leclourec P (1995). Infrared spectroscopic study of uranyl biosorption by fungal biomass and materials of biological origin. *Environ. Sci. Technol.* 29: 2496-2503.
- Gulati R, Saxena RK, Gupta R (1999). Fermentation waste of *Aspergillus terreus*: Promising copper bio-indicator. *Curr. Sci.* 77: 1359-1360.
- Gupta R, Ahuja P, Khan S, Saxena RK, Mohapatra H (2000). Microbial biosorbents: Meeting challenges of heavy metal pollution in aqueous solutions. *Curr. Sci.* 78: 967-973.
- Huang CP, Huang CP (1996). Application of *Aspergillus oryzae* and *Rhizopus oryzae*. *Wat. Res.* 30: 1985-1990.
- Jacob CT, Azariah J, Hilda A, Gopinath S (1998). Decolorization of procion red MX-5B and coloured textile effluent using *Phanerochaete chrysosporium*. *J. Environ. Biol.* 19: 259-264.
- Jimenez AM, Borja R (1997). Influence of aerobic pretreatment with *Penicillium decumbens* on the anaerobic digestion of beet molasses alcoholic fermentation wastewater in suspended and immobilized cell bioreactors. *J. Chem. Technol. Biotechnol.* 69: 193-202.
- Jin B, van Leeuwen J, Yu Q, Patel B (1998). Utilisation of starch processing wastewater for production of microbial biomass protein and alpha-amylase by *Aspergillus oryzae*. *Bioresource Technol.* 66: 201-206.
- Jin B, van Leeuwen J, Patel B (1999a). Production of fungal protein and glucoamylase by *Rhizopus oligosporus* from starch processing wastewater. *Process Biochem.* 34: 59-65.
- Jin B, van Leeuwen J, Yu Q, Patel B (1999b). Mycelial morphology and fungal protein production from starch processing wastewater in submerged cultures of *Aspergillus oryzae*. *Process Biochem.* 34: 335-340.
- Jin B, van Leeuwen J, Yu Q, Patel B (1999c). Screening and selection of microfungi for microbial biomass protein production and water reclamation from starch processing wastewater. *J. Chem. Technol. Biotechnol.* 74: 106-110.
- Jin B, Yu Q, van Leeuwen J (2001). A bioprocessing mode for simultaneous fungal biomass protein production and wastewater treatment using an external air-lift bioreactor. *J. Chem. Technol. Biotechnol.* 76: 1041-1048.
- Kahmark KA, Unwin JP (1999). Pulp and paper effluent management. *Water Environ. Res.* 71: 836-852.
- Kapoor A, Viraraghavan T (1995). Fungal biosorption: an alternative treatment option for heavy metal bearing wastewater. A review. *Bioresource Technol.* 53: 196-206.
- Kapoor A, Viraraghavan T (1997). Heavy metals biosorption site in *Aspergillus niger*. *Bioresource Technol.* 61: 221-227.
- Kapoor A, Viraraghavan T (1998a). Removal of heavy metals from aqueous solution using immobilized fungal biomass in continuous mode. *Wat. Res.* 32: 1968-1977.
- Kapoor A, Viraraghavan T (1998b). Application of immobilized *Aspergillus niger* biomass in the removal of heavy metals from an industrial wastewater. *J. Environ. Sci. Health.* 33: 1507-1514.
- Kapoor A, Viraraghavan T, Cullimore DR (1999). Removal of heavy metals using the fungus *Aspergillus niger*. *Bioresource Technol.* 70: 95-104.
- Karavaiko GI, Zakharova VI, Avakyan ZA, Strizhko LS (1996). Selective extraction of noble metals from solutions by microorganisms. *Appl. Biochem. Microbiol.* 32: 507-510.
- Kida K, Morimura S, Abe N, Sonoda Y (1995). Biological treatment of shochu distillery wastewater. *Process Biochem.* 30: 125-132.
- Kirby N, Marchant R, McMullan G (2000). Decolorisation of synthetic textile dyes by *Phebia tremellosa*. *FEMS Microbiol. Lett.* 188: 93-96.
- Kissi M, Mountadar M, Assobhei O, Gargiulo E, Palmieri G, Giardina P, Sannia G (2001). Roles of two white-rot basidiomycete fungus in

- decolourisation and detoxification of olive mill waste water. Appl. Microbiol. Biotechnol. 57: 221-226.
- Knapp JS, Newby PS (1999). The decolorization of chemical industry effluent by white rot fungi. Wat. Res. 33: 575-577.
- Kramer M, Meisch HU (1999). New metal-binding ethyldiamino and dicarboxy-products from *Aspergillus niger* industrial wastes. Biometals 12: 241-246.
- Kratochvil D, Volesky B (1998). Advances in biosorption of heavy metals. Trends Biotechnol. 16: 291-300.
- Kumar V, Wati L, Nigam P., Yadav BS, Singh D, Marchant R (1998). Decolorization and biodegradation of anaerobically digested sugarcane molasses spent wash effluent from biomethanation plants by white-rot fungi. Process Biochem. 33: 83-88.
- Lal DN (1980). The influence of some ammonium compounds on the production of citric acid by *Aspergillus niger* AL. 29. Ind. J. Agri. Chem. 13: 153-157
- Lilly VM, Barnett HL (1951). Physiology of the fungi. McGraw-Hill Book Co., 1<sup>st</sup> ed. New York, New York.
- Livernoche D, Jurasek L, Desrochers M, Dorica J (1983). Removal of color from kraft mill wastewaters with cultures of white-rotfungi and with immobilized mycelium of *Coriolus versicolor*. Biotechnol. Bioeng. 225: 2055-2065.
- Manzanares P, Fajardo S, Martin C (1995). Production of ligninolytic activities when treating paper pulp effluents by *Trametes versicolor*. J. Biotechnol. 43: 125-132.
- Marwaha S, Grover R, Prakash C, Kennedy J (1998). Continuous bioleaching of black liquor from the pulp and paper industry using immobilized cell system. J. Chem. Technol. Biotechnol. 73: 292.
- Mashitah MD, Zulfadhly Z, Bhatia S (1999). Binding mechanism of heavy metals biosorption by *Pycnoporus sanguineus*. Artif. Cell. Blood Substit. Immobil. Biotechnol. 27: 441-445.
- Matheickal JT, Yu Q (1997). Biosorption of lead (II) from aqueous solutions by *Phellinus badius*. Miner. Eng. 10: 947-957.
- McHale AP, McHale S (1994). Microbial biosorption of metals. Potential in the treatment of metal pollution. Biotechnol. Adv. 12: 647-652.
- McKay G, Ho YS, Ng JCY (1999). Biosorption of copper from waste water : A review. Separ. Purif. Method. 28: 87-125.
- McMullan G, Meehan C, Conneely A, Kirby N, Robinson T, Nigam P, Banat IM, Marchant R, Smyth WF (2001). Microbial decolorisation and degradation of textile dyes. Appl. Microbiol. Biotechnol. 56: 81-87.
- Messner K, Ertler G, Jaklin-Farcher S, Boskosvsky P, Regensberger U, Blada A (1990). Treatment of bleach plant effluent by the mycopor system. In *biotechnology in Pulp and Paper Manufacture, Applications and Fundamental investigations*. Eds T. K. Kirk, H.-M. Chang, Butterworth-Heinemann, Boston, pp. 245-251.
- Mester T, Tien M (2000). Oxidation mechanism of ligninolytic enzymes involved in the degradation of environmental pollutants. Int. Biodeterior. Biodegrad. 46: 51-59
- Meyer A, Wallis FM (1997). The use of *Aspergillus niger* (strain 4) biomass for lead uptake from aqueous systems. Water S. A. 23: 187-192.
- Miranda MP, Benito GG, San Cristobal N, Nieto CH (1996). Color elimination from molasses wastewater by *Aspergillus niger*. Bioresource Technol. 57: 229-235.
- Miyata N, Mori T, Iwahori K, Fujita M (2000). Microbial decolorization of melanoidin containing wastewaters: combined use of activated sludge and the fungus *Coriolus hirsutus*. J. Biosci. Bioeng. 89: 145-150.
- Modak JM, Natarajan KA (1995). Biosorption of metals using nonliving biomass. A review. Minerals Metall. Process. 12: 189-196.
- Modak JM, Natarajan KA, Saha B (1996). Biosorption of copper using waste *Aspergillus niger* biomass. Minerals Metall. Process. 12: 189-196.
- Mogollon L, Rodriguez R, Larrota W, Ramirez N, Torres R (1998). Biosorption of nickel using filamentous fungi. Appl. Biochem. Biotech. 70: 593-601.
- Morimura S, Kida K, Sonoda Y (1992). The influence of shear stress and its reduction in the production of saccharifying enzyme from shochu distillery wastewater by *Aspergillus awamori* var kawachi. J. Ferment. Bioeng. 78: 160-163.
- Morimura S, Kida K, Nakagawa M, Sonoda Y (1994a). Production of protease using wastewater from the manufacture of shochu. J. Ferment. Bioeng. 77: 183-187.
- Morimura S, Kida K, Nakagawa M, Sonoda Y (1994b). Production of fungal proteins by *Aspergillus awamori* var kawachi grown in shochu distillery wastewater. J. Ferment. Bioeng. 78: 160-163.
- Mullen MD, Wolf DC, Beveridge TJ, Bailey GW (1992). Sorption of heavy metals by soil fungi *Aspergillus niger* and *Mucor rouxii*. Soil Biol. Biochem. 24: 129-135.
- Murado MA, Gonzalez MP, Pastrana L, Siso MIG, Miron J, Montemayor MI (1993). Enhancement of the bioproduction potential of an amylaceous effluent. Bioresource Technol. 44: 155-163.
- Nagarathnamma R, Bajpai P (1999). Decolorization and detoxification of extraction stage effluent from chlorine bleaching of kraft pulp by *Rhizopus oryzae*. Appl. Environ. Microbiol. 65: 1078-1082.
- Nagarathnamma R, Bajpai P, Bajpai PK (1999). Studies on decolorization and detoxification of chlorinated lignin compounds in kraft bleaching effluents by *Ceriporiopsis subvermispora*. Process Biochem. 34: 939-948.
- Nieto LM, Cormenzana AR, Pareja PG, Hoyos SEG (1992). Phenolic compounds biodegradation of olive mill wastewater with *Aspergillus terreus*. Grasas y Aceites. 43: 75-81.
- Nigam P, Banat IM, Oxspring D, Marchant R, Singh R, Smyth WF (1995). A new facultative anaerobic filamentous fungus capable of growth on recalcitrant textile dyes as sole carbon source. Microbios 84: 171-185.
- Niyogi S, Abraham TE, Ramakrishna SV (1998). Removal of chromium (VI) ions from industrial effluents by immobilized biomass of *Rhizopus arrhizus*. J. Sci. Ind. Res. India 57: 809-816.
- Ohmomo S, Kainuma M, Sirianuntapiboon S, Aoshima I, Atthasampunna P (1988). Adsorption of melanoidin to the mycelia of *Aspergillus oryzae* Y-2-32. Agric. Biol. Chem. 52: 381-386.
- Ollikka P, Harjunpaa T, Palmu K, Mantsala P, Suominen I (1998). Oxidation of Crocein Orange G by lignin peroxidase isoenzymes. Kinetics and effect of H<sub>2</sub>O<sub>2</sub>. Appl. Biochem. Biotechnol. 75: 307-321.
- O'Neill C, Hawkes FR, Hawkes DL, Lourenço ND, Pinheiro HM, Delée W (1999). Colour in textile effluents-sources, measurement, discharge contents and simulation: a review. J. Chem. Technol. Biotechnol. 74: 1009 -1018.
- Palma C, Moreira MT, Mielgo I, Feijoo G, Lema JM (1999). Use of a fungal bioreactor as a post treatment step for continuous decolorisation of dyes. Wat. Sci. Technol. 40: 131-136.
- Payman MA, Mahnaz MA (1998). Decolorization of textile effluent by *Aspergillus niger* (marine and terrestrial). Fresen. Environ. Bull. 7: 1-7.
- Perez J, Saez L, De La Rubia T, Martinez J (1997). *Phanerochaete flavidio-alba* ligninolytic activities and decolorization of partially biodepurated paper millwastes. Wat. Res. 31: 495-502.
- Podgornik H, Grgic I, Perdih A (1999). Decolorization rate of dyes using lignin peroxidases of *Phanerochaete chrysosporium*. Chemosphere 38: 1353-1359.
- Pointing SB, Vrijmoed LLP (2000). Decolorization of azo and triphenylmethane dyes by *Pycnoporus sanguineus* producing laccase as the sole phenoloxidase. World J. Microbiol. Biotechnol. 16: 317-318.
- Polman JK, Breckenridge CR (1996). Biomass-mediated binding and recovery of textile dyes from waste effluents. Textil chemist and colorist. 28: 31-35.
- Prasertsan P, Kittikul AH, Maneesri J, Oi S (1997). Optimization for xylanase and cellulase production from *Aspergillus niger* ATCC 6275 in palm oil mill wastes and its application. World J. Microbiol. Biotechnol. 13: 555-559.
- Price MS, Classen JJ, Payne GA (2001). *Aspergillus niger* absorbs copper and zinc from swine wastewater. Bioresource Technol. 77: 41-49.
- Radzio R, Kuck U (1997). Synthesis of biotechnologically relevant heterologous proteins in filamentous fungi. Process Biochem. 32: 529-539.
- Raghukumar C, Chandramohan D, Michel FC Jr, Reddy CA (1996). Degardation of lignin and decolorization of paper mill bleach plant effluent by marine. Biotechnol. Lett. 18: 105-106.
- Raghukumar C, Rivonkar G (2001). Decolorization of molasses spent wash by white-rot fungus *Flavodon flavus*, isolated from a marine

- habitat. *Appl. Microbiol. Biotechnol.* 55: 510-514.
- Reyes P, Pickard MA, Vazquez-Duhalt R (1999). Hydroxybenzotriazole increases the range of textile dyes decolorized by immobilized laccase. *Biotechnol. Lett.* 21: 875-880.
- Robles A, Lucas R, de Cienfuegos GA, Galvez A (2000). Biomass production and detoxification of wastewaters from the olive oil industry by strains of *Penicillium* isolated from wastewater disposal ponds. *Bioresource Technol.* 74: 217-221.
- Robinson T, McMullan G, Marchant R, Nigam P (2001). Remediation of dyes in textile effluent: a critical review on current treatment technologies with a proposed alternative. *Bioresource Technol.* 77: 274-255.
- Rodriguez E, Pickard MA, Vazquez-Duhalt R (1999). Industrial dye decolorization by laccases from lignolytic fungi. *Curr. Microbiol.* 38: 27-32.
- Sag Y, Acikel U, Aksu Z, Kutsal T (1998). A comparative study for the simultaneous biosorption of Cr(VI) and Fe(III) on *C. vulgaris* and *R. arrhizus*: application of the competitive adsorption models. *Process Biochem.* 33: 273-281.
- Sarret G, Manceau A, Spadini L, Roux JC, Hazemann JL, Soldo Y, Eybert-Berard L, Menthonnex JJ (1998). Structural determination of Zn and Pb binding sites in *Penicillium chrysogenum* cell walls by EXAFS spectroscopy. *Environ. Sci. Technol.* 11: 1648-1655.
- Sarret G, Manceau A, Spadini L, Roux JC, Hazemann JL, Soldo Y, Eybert-Berard L, Menthonnex JJ (1999). Structural determination of Pb binding sites in *Penicillium chrysogenum* cell walls by EXAFS spectroscopy and solution chemistry. *J. Synchrotron Radiat.* 6: 414-416.
- Savvaidis I (1998). Recovery of gold from thiourea solutions using microorganisms. *Biometals* 11: 145-151.
- Sayadi S, Ellouz R (1995). Roles of lignin peroxidase and manganese peroxidase from *Phanerochaete chrysosporium* in the decolorization of olive mill wastewaters. *Appl. Environ. Microbiol.* 61: 1098-1103.
- Sayadi S, Allouche N, Jaoua M, Aloui F (2000). Detrimental effects of high molecular mass polyphenols on olive mill wastewater biotreatment. *Process Biochem.* 35: 725-735.
- Setti L, Maly S, Iacondini A, Spinuzzi G, Pifferi PG (1998). Biological treatment of olive milling waste waters by *Pleurotus ostreatus*. *Annali Chimica.* 88: 201-210.
- Shaul GM, Dempsey CR, Dostal KA, Lieberman RJ (1985). Fate of azo dyes in the activated sludge process. In *Proceedings 41 st Purdue University Industrial Waste Conference*, pp 603-611.
- Sing C, Yu J (1998). Copper adsorption and removal from water by living mycelium of white-rot fungus *Phanerochaete chrysosporium*. *Wat. Res.* 32: 2746-2752.
- Soares CHL, Duran N (1998). Degradation of low and high molecular mass fraction of kraft E1 effluent by *Trametes villosa*. *Environ. Technol.* 19: 883-891.
- Sumathi S, Phatak V (1999). Fungal treatment of bagasse based pulp and paper mill wastes. *Environ. Technol.* 20: 93-98.
- Swamy J, Ramsay JA (1999). Effect of Mn<sup>2+</sup> and NH<sub>4</sub><sup>+</sup> concentration on laccase and manganese peroxidase production and Amaranth decoloration by *Trametes versicolor*. *Appl. Microbiol. Biotechnol.* 51: 391-396.
- Tatarko M, Bumpus JA (1998). Biodegradation of Congo Red by *Phanerochaete chrysosporium*. *Wat. Res.* 32: 1713-171.
- Tereshina VM, Mar'in AP, Kosyakov NV, Kozlov VP, Feofilova EP (1999). Different metal sorption capacities of cell wall polysaccharides of *Aspergillus niger*. *Appl. Biochem. Microbiol.* 35: 389-392.
- Thanh NC, Simar RE (1973). Biological treatment of domestic sewage by fungi. *Mycopathol. Mycol. Applicata* 51: 223-232.
- Thomke S, Rundgren M (1980). Nutritional evaluation of the white-rot fungus *Sporotrichum pulverulentum* as a feedstuff to rats, pigs, and sheep. *Biotechnol. Bioeng.* 22: 2285-2303.
- Tobin JM, Roux JC (1998). Mucor biosorbent for chromium removal from tanning effluent. *Wat. Res.* 32, 1407-1416.
- Tsezos M, Volesky B (1982). The mechanism of thorium biosorption by *Rhizopus arrhizus*. *Biotechnol. Bioeng.* 24: 955-969.
- Vaidya AA, Datye KV (1982). Environmental pollution during chemical processing of synthetic fibers. *Colourate* 14: 3-10.
- van Driessel B, Christov L (2002). Adsorption of colour from a bleach plant effluent using biomass and cell wall fractions from *Rhizomucor pusillus*. *J. Chem. Technol. Biotechnol.* 77: 155-158.
- Vandevivere PC, Bianchi R, Verstraete W (1998). Treatment and reuse of wastewater from the textile wet-processing industry: Review of emerging technologies. *J. Chem. Technol. Biotechnol.* 72: 289-302.
- Vassilev N, Fenice M, Federici F, Azcon R (1997). Olive mill wastewater treatment by immobilized cells of *Aspergillus niger* and its enrichment with soluble phosphate. *Process Biochem.* 32: 617-620.
- Vassilev N, Vassileva M, Azcon R, Fenice M, Barea JM (1998). Fertilizing effect of microbially treated olive mill wastewater on *Trifolium* plants. *Bioresource Technol.* 66: 133-137.
- Veglio F, Beolchini F (1997). Removal of metals by biosorption : a review. *Hydrometallurgy.* 44: 301-316.
- Vinciguerra V, Dannibale A, Dellemonache G, Sermanni GG (1995). Correlated effects during the bioconversion of waste olive waters by *Lentinus edodes*. *Bioresource Technol.* 51: 221-226.
- Volesky B (1994). Advances in biosorption of metals: Selection of biomass types. *FEMS Microbiol. Rev.* 14: 291-302.
- Volesky B, Holan ZR (1995). Biosorption of heavy metals. *Biotechnol. Progr.* 11: 235-250.
- Wesenberg D, Buchon F, Agathos SN (2002). Degradation of dye-containing textile effluent by the agaric white-rot fungus *Clitocybula duseinii*. *Biotechnol. Lett.* 24: 989-993.
- Wesenberg D, Kyriakides I, Agathos SN (2003). White-rot fungi and their enzymes for the treatment of industrial dye effluents. *Biotechnol. Adv.* 22: 161-187.
- Wong YX, Yu J (1999). Laccase catalyzed decolorization of synthetic dyes. *Wat. Res.* 33: 3512-3520.
- Wongwicharn A, Harvey LM, McNeil B (1999). Secretion of heterologous and native proteins, growth and morphology in batch cultures of *Aspergillus niger* B1-D at varying agitation rates. *J. Chem. Technol. Biotechnol.* 74: 821-828.
- Xu JF, Wang LP, Ridgway D, Gu TY, Moo-Young M (2000). Increased heterologous protein production in *Aspergillus niger* fermentation through extracellular proteases inhibition by pellet growth. *Biotechnol. Prog.* 16: 222-227.
- Yan GY, Viraraghavan T (2000). Effect of pretreatment on the biosorption of heavy metals on *Mucor rouxii*. *Water S. A.* 26: 119-123.
- Yang F-C, Lin I-H (1998). Production of acid protease using thin stillage from a rice -spirit distillery by *Aspergillus niger*. *Enzyme Microb. Technol.* 23: 397-402.
- Yesilada O, Fiskin K, Yesilada E (1995). The use of white rot fungus *Funalia trogii* (Malatya) for the decolorization and phenol removal from olive mill wastewater. *Environ. Technol.* 16: 95-100.
- Yesilada O, Sik S, Sam M (1998). Biodegradation of olive oil mill wastewater by *Coriolus versicolor* and *Funalia trogii*: effects of agitation, initial COD concentration, inoculum size and immobilization. *World J. Microbiol. Biotechnol.* 14: 37-42.
- Yin PH, Yu QM, Ling Z (1999). Biosorption removal of cadmium from aqueous solution by using pretreated fungal biomass cultured from starch wastewater. *Wat. Res.* 33: 1960-1963.
- Yokoi H, Aratake T, Nishio S, Hirose J, Hayashi J, Takasaki Y (1998). Chitosan production from Shochu distillery wastewater by fungi. *J. Ferment. Bioeng.* 85: 246-249.
- Young L, Yu J (1997) Ligninase- catalysed decolorization of synthetic dyes. *Wat. Res.* 31: 1187-1193.
- Yu QM, Kaewsarn P (1999). A model for pH dependent equilibrium of heavy metal biosorption. *Korean J. Chem. Eng.* 16: 753-757.
- Zhang L, Zhao L, Yu YT, Chen CZ (1998). Removal of lead from aqueous solution by non-living *Rhizopus nigricans*. *Wat. Res.* 32: 1437-1444.
- Zheng Z, Levin RE, Pinkham JL, Shetty K (1999). Decolorization of polymeric dyes by a novel *Penicillium* isolate. *Process Biochem.* 34: 31-37.
- Zhou JL (1999). Zn biosorption by *Rhizopus arrhizus* and other fungi. *Appl. Microbiol. Biotechnol.* 51: 686-693.